

EXHIBIT A



SHORT REPORT

A novel type of mutation in the cysteine rich domain of the RET receptor causes ligand independent activation

DH Arlt¹, B Baur¹, B Wagner² and W Höppner^{*1}

¹Institute for Hormone and Fertility Research, University of Hamburg, Grandweg 64, 22529 Hamburg, Germany; ²University Hospital Eppendorf, Department of Neurosurgery, 20246 Hamburg, Germany

Multiple endocrine neoplasia type 2A (MEN 2A) is a dominantly inherited cancer syndrome, which involves the triad of MTC, pheochromocytoma, and hyperparathyroidism. Missense mutations in one of six cysteine codons in the extracellular cysteine-rich domain of the *RET* proto-oncogene predispose to this disease. These mutations cause ligand-independent constitutive activation of the tyrosine kinase receptor by the formation of disulfide-bonded homodimers. We examined a different type of mutation, which results in an additional cysteine in the cysteine rich domain. A duplication of 9 bp in the first case resulted in an insertion of three amino acids between codon 633 and 634. In the second case a 12 bp duplication in exon 11 results in four additional amino acids between codon 634 and 635. Here we demonstrate that an additional cysteine causes a ligand independent dimerization of the RET receptor in transfected NIH3T3 cells, which results in an activation of the intracellular tyrosine kinase. *Oncogene* (2000) 19, 3445–3448.

Keywords: MEN 2; *RET*-proto-oncogene; additional cysteine 634; ligand independent dimerization

The *RET* proto-oncogene codes for a receptor tyrosine kinase which plays a role in the development of the neural crest and its derivatives. Besides this, RET is important for kidney development (Pachnis *et al.*, 1993; Avantsgiato *et al.*, 1994). In adult humans RET is expressed in cells of the nervous system, in C-cells, the adrenal medulla and the parathyroids (Nakamura *et al.*, 1994). It has been shown that the glial cell line-derived neurotrophic factor (GDNF), neurturin (NTN) and persephin are functional ligands for the RET receptor (Durbec *et al.*, 1996; Baloh *et al.*, 1997; Milbrandt *et al.*, 1998). Activation of the tyrosine kinase is mediated by glycosylphosphatidylinositol-(GPI)-anchored cell surface RET-coreceptors GDNFR- α 1–3 (Treanor *et al.*, 1996; Jing *et al.*, 1996; Baloh *et al.*, 1997).

Mutations in the *RET* proto-oncogene are involved in several inherited and non inherited diseases. Inactivating mutations cause familial and sporadic Hirschsprung disease, a syndrome of congenital absence of enteric innervation (Romeo *et al.*, 1994). Activating mutations have been identified as the underlying cause of multiple endocrine neoplasia Type 2 (MEN 2) and familial medullary thyroid carcinoma (FMTC) (Donis-Keller *et al.*, 1993; Mulligan *et al.*,

1993). Mutations in one of six cysteine codons in exon 10 or exon 11 occur in most cases of MEN 2A and FMTC. These mutations convert *RET* into a dominant transforming oncogene (Santoro *et al.*, 1995). Substitutions of cysteine residues related to the extracellular cysteine-rich domain result in ligand independent dimerization of the receptor and constitutive activation by autophosphorylation of specific tyrosine-residues in the intracellular kinase domain (TK) (Santoro *et al.*, 1995). Probably the Cys residues, disrupted by the MEN 2A mutations, are normally involved in intramolecular disulfide bonds. The mutations may render the partner Cys available for aberrant disulfide binding with other mutant RET or wt-RET molecules and thus form activated dimers. The activated RET binds the adaptor proteins Shc, Grb2 and PLC- γ (Borrello *et al.*, 1994, 1996). Rare MEN 2A/FMTC-associated mutations involve the intracellular kinase domain. In particular the codons 790, 791 and 804 in the TK1 domain are affected (Berndt *et al.*, 1998; Fattoruso *et al.*, 1998).

Recently we described a novel type of MEN 2A associated mutation which results in an additional Cys 634 in exon 11. In the germline of a patient with the complete phenotype of MEN 2A a duplication of 9 bp representing the codons 634, 635 and 636 was found (Höppner *et al.*, 1998). In the second case a 12 bp-duplication which results in additional four amino acids between codon 634 and 635 was detected in a family with 14 affected members (Höppner *et al.*, 1997). While the 9 bp-duplication is in frame, the inserted 12 bp sequence is not in frame which results in an additional histidine residue at the 5' end of the breakpoint. The 12 bp duplication is associated with a different MEN 2A phenotype. Besides MTC an unusual accumulation of pHPT occurs, none of the 14 affected family members have pheochromocytomas (Höppner *et al.*, 1997).

To examine the mechanism of activation of the mutated RET, we cloned the *RET*-cDNA of the short isoform into the eucaryotic expression vector pRC/CMV2 and inserted the mutations described above by fragmental cloning in the case of the 9 bp-duplication (+9 bp-RET) and by PCR-mutagenesis for the 12 bp-duplication (+12 bp-RET). The mouse fibroblast cell line NIH3T3 was transfected and clones expressing wt-RET, +9 bp-RET and +12 bp-RET were selected. TT-cells, a human medullary thyroid carcinoma cell line which is heterozygous for one of the classical mutations in codon 634 (C634W) of the *RET* proto-oncogene was also cultured. After cell lysis we analysed the proteins by nonreducing and reducing SDS-polyacrylamid gel electrophoresis (Figure 1).

*Correspondence: W Höppner

Received 9 May 2000; revised 12 May 2000; accepted 16 May 2000

Ligand independent dimerization of the +9 bp- and +12 bp-RET

In agreement with previous reports (Santoro *et al.*, 1995) the mutation C634W results in ligand independent dimerization of the RET receptor, which is represented by the 300 kDa band of RET expressed in TT-cells (Figure 1a, lane 1). The monomeric forms migrate as 150 and 170 kDa bands. The +9 bp-RET (Figure 1a, lane 2) and +12 bp-RET (Figure 1a, lane 3) also migrate as dimers whereas the wt-expressing cells only show the monomeric RET (Figure 1a, lane 4). Therefore it can be concluded that the additional Cys 634 results in a ligand independent dimerization of the RET receptor.

Formation of RET-dimers by intermolecular disulfide linkages

As shown in previous studies (Santoro *et al.*, 1995), under reducing conditions the C634W-RET migrates as a monomer (Figure 1b, lane 1). This serves as an evidence that the mutation causes a ligand independent




Figure 1 (a) Western blot analysis of wt and mutated RET receptors under nonreducing conditions. Comparable amounts of total cellular proteins from TT-cells (lane 1), and NIH3T3 cells stable transfected with +9 bp-RET (lane 2), +12 bp-RET (lane 3) and wt-RET (lane 4) were dissolved in loading buffer without DTT and subjected to SDS-PAGE. The proteins were detected after Western blotting with a polyclonal anti-RET antibody (Santa Cruz). Immunocomplexes were visualized by enhanced chemiluminescence detection kit (ECL, Pierce) using anti-goat coupled to horseradish peroxidase (DIANOVA). +9 bp-RET, +12 bp-RET and C634W-RET (lane 1) migrate as dimeric forms represented by the band at 300 kDa while the wt-RET migrates only as monomeric forms at 150 and 170 kDa. (b) Under reducing conditions +9 bp-RET gene product (lane 2) and the +12 bp protein (lane 3) show a molecular weight of 150 and 170 kDa. The positive control, C634W-RET (lane 1) also migrates as monomers

dimerization by formation of intermolecular disulfide linkages. Under reducing conditions also the +9 bp-RET (Figure 1b, lane 2) and the +12 bp gene product (Figure 1b, lane 3) are only detectable as monomeric forms. Obviously the additional Cys 634 also causes interchain disulfide linkages which result in ligand independent dimerization.

Constitutive activation of the +9 bp- and +12 bp-RET tyrosine kinase in NIH3T3 cells

To investigate the ligand independent activation of the mutated receptors, RET was immunoprecipitated from the various NIH3T3 transfectants and assayed for phosphotyrosine (pTyr) content, a hallmark of receptor activation. In case of the TT cells we could confirm previous results (Santoro *et al.*, 1995) which show that the mutation C634W causes an autophosphorylation of the intracellular tyrosine kinase (Figure 2b, lane 1). This also has been shown for RET receptors with Cys 634 substitution expressed in NIH3T3 cells. Our results demonstrate that the +9 bp- and +12 bp-RET proteins also are autophosphorylated (Figure 2b, lane 2 and lane 3). In comparison with the TT-cells there is a similar amount of pTyr in the mutated receptors with the additional cysteine 634. These results show that this type of mutation causes a constitutive activation of the receptor tyrosine kinase after dimerization.

One could speculate that overexpression of the wt-RET is sufficient to trigger the kinase activity. To determine whether overexpression or the additional cysteine 634 causes the constitutive activation, we compared cells with less expression of the +12 bp-




Figure 2 Activation of the tyrosine kinase of the RET receptor. Comparable amounts of total cellular proteins from the indicated transfectants were immunoprecipitated with a polyclonal anti-RET antibody. The immunoprecipitates were then either immunoblotted with anti-RET (a) or a monoclonal antibody to phosphotyrosine (pTyr PY99) (Santa Cruz) (b). In case of the duplication in exon 11 the RET product shows high amounts of pTyr (b: lanes 2 and 3). The RET protein of the TT-cells (a: lane 1) shows a low signal for pTyr (b: lane 1) but in relation to the expression level it is comparable with the +9 bp-RET and the +12 bp-RET

RET and cells with a higher expression of wt-RET. The wt-RET displays autophosphorylation but in comparison with the +12 bp-RET the amount of pTyr is significantly less (data not shown). This demonstrates that it is not the level of expression that causes ligand independent activation but the mutation itself.

Grb2-binding by the activated RET

It has been shown that activated RET receptors bind Grb2 and trigger intracellular signal pathways. To confirm that receptor activation mediated by an additional Cys 634 leads to Grb2 binding we performed coprecipitation experiments of Grb2 and the RET protein. Our results show that the receptors with the duplications in exon 11 bind Grb2 (Figure 3b, lane 2 and lane 3) as is the case for the RET protein with the classical mutation in codon 634 (Figure 3b, lane 1). This is also the case for the +12 bp-RET clones with low expression (Figure 3b, lane 3).

These data present evidence for receptor activation by a duplication of a sequence including the critical cysteine residue in codon 634. *In vitro* no differences in the mechanism of activation occur between the 9 bp- and the 12 bp-duplication. The phenotype of the +12 bp-transfectants differs from that of the +9 bp-transfectants (data not shown). The latter show the complete transformed phenotype with cell clusters and spindle shape whereas the +12 bp-transfectants only express cell clusters but exhibit the same cell shape as wt-transfected cells. Interestingly the *in vivo* phenotype of the +12 bp-family differs from that of families with

the classical mutations and the +9 bp-patient. The fact that no pheochromocytoma occurs but that pPHT does suggest that tissue specific mechanisms exist.

The different mutations investigated in this study alter significantly the chemical properties of the cysteine-rich domain in close vicinity to the plasma membrane and lead to considerably different spacing of the cysteine codons. In contrast to the missense mutation in codon 634 the two duplication mutations leave the first extracellular cysteine at the same distance from the cell surface as in the wt-RET (Figure 4). Also the recently published deletion of six base pairs (Bongarzone *et al.*, 1999), which represents an activating mutation too, still leaves the cysteine 634 in place (Figure 4). Thus the distance of the cysteine relative to the plasma membrane seems not to be essential.

It has not been shown whether Cys 634 and Cys 630 are involved in intrachain disulfide bonds *in vivo*. The fact that substitutions and insertions of Cys 634, just as deletions between Cys 634 and Cys 630, result in ligand independent dimerization suggest an essential function of this domain in transition from inactive monomers to activated dimers. The distance between Cys 634 and the next extracellular cysteine in codon 630 is two amino acids in the Δ6-RET, three amino acids in the +9 bp-RET and four amino acids in the +12 bp-RET and the wt-RET receptor respectively (Figure 4). If intrachain disulfide bonds should form between these cysteines, the energy of the bonds may differ significantly or may even become sterically impossible. In the case of cysteine duplication, there may be a competition between disulfide linkage of the new cysteine with cysteine 634 and 630. For the 12 bp duplication, the distance in both directions is the same and equals the wild type conditions. The only difference is an Asp-His exchange in the three amino acids between the cysteines. In this case disulfide linkages may be equally likely to both sides. Thus




Figure 3 Grb2-binding of the mutated RET. Comparable amounts of total cellular proteins from the indicated transfectants were immunoprecipitated with an anti-RET antibody. The immunoprecipitates were then either immunoblotted with anti-RET (a) or antiGrb2 (Dianova) (b). Although the +12 bp-RET shows a low expression (a: lane 3) in comparison with the RET protein expressed in TT-cells (a: lane 1) and the +9 bp-RET (a: lane 2), the amount of coprecipitated Grb2 is comparable (b: lane 3)




Figure 4 Schematic illustration of the amino acid sequence in the vicinity of the plasma membrane. The wt-sequence shows the original arrangement of Cys 630 and Cys 634. Four types of mutation are described which affect this domain. The classical mutation is a substitution of Cys 634. In this figure the mutation Cys 634Arg is shown, but other amino acid changes are described. Another type of mutation is a 6 bp-deletion (Δ6) which does not affect Cys 634 directly, but moves Cys 630 and Cys 634 closer to one another. The type of mutation described in this paper is a duplication of Cys 634 (+9 bp and +12 bp). In the case of the 9 bp-duplication, the sequence Thr-Arg-Cys is inserted (box), the 12 bp-duplication results in the additional sequence Cys-Leu-Glu-His (box)

either cysteine 630 or 634 may be available for interchain disulfide linkage.

For the 9 bp duplication the distance to cysteine 630 is four codons and to cysteine 634 is three codons. The disulfide linkage between cysteine 630 and the new cysteine would be the same as in the wild type and may be preferred, one cysteine remaining free and this may become involved in interchain disulfide linkage. For the $\Delta 6$ bp-*RET* the distance of the two cysteines may be too small to facilitate intrachain disulfide linkage (Bongarzone et al., 1999).

It is not certain whether intrachain disulfide bonds occur between Cys 630 and Cys 634. Another possibility is that these residues form bridges with more distant cysteine residues. In this case it is possible that displacement occurs between the inserted cysteine and the original Cys 634. The different pattern of disulfide bonds would alter the stability of interchain disulfide bridges for the 12 bp-duplication.

References

- Avantaggiato V, Dathan NA, Grieco M, Fabien N, Lazzaro D, Fusco A, Simeone A and Santoro M. (1994). *Cell Growth Differ.*, **5**, 305–311.
- Baloh RH, Tansey MG, Golden JP, Creedon DJ, Heuckeroth RO, Keck CL, Zimonjic DB, Popescu NC, Johnson EM and Milbrandt J. (1997). *Neuron*, **18**, 793–802.
- Berndt I, Reuter M, Saller B, Frank-Raue K, Groth P, Grußendorf M, Raue F, Ritter MM and Höppner W. (1998). *J. Clin. Endocr. Metab.*, **83**, 770–774.
- Bongarzone I, Vigano E, Alberti L, Mondellini P, Uggeri M, Pasini B, Borrello MG and Pierotti MA. (1999). *Oncogene*, **34**, 4833–4838.
- Borrello MG, Pelicci G, Arighi E, De Filippis L, Bongarzone I, Rizetti MG, Pellici PG and Pierotti MA. (1994). *Oncogene*, **9**, 1661–1668.
- Borrello MG, Alberti L, Arighi E, Bongarzone I, Battistini C, Bardelli A, Pasini B, Piutti C, Rizetti MG, Mondellini P, Radice MT and Pierotti MA. (1996). *Mol. Cell Biol.*, **16**, 2151–2163.
- Donis-Keller H, Dou S, Chi D, Carlson KM, Toshima K, Lairmore TC, Howe JR, Moley JF, Goodfellow P and Wells Jr SA. (1993). *Hum. Mol. Genet.*, **2**, 851–856.
- Durbec P, Marcos-Gutierrez CV, Kilkenny M, Grigoriou M, Wartiovaara K, Suvanto P, Smith D, Ponder B, Constantini F, Saarma M, Sariola H and Pachnis V. (1996). *Nature*, **381**, 789–793.
- Fattoruso O, Quadro L, Libroia A, Verga U, Lupoli G, Cascone E and Colantuoni V. (1998). *Hum. Mutat.*, (Suppl 1), 167–171.
- Höppner W and Ritter MM. (1997). *Hum. Mol. Genet.*, **6**, 587–590.
- Höppner W, Dralle H and Brabant G. (1998). *Hum. Mutat.*, (Suppl 1), 128–130.
- Jing S, Wen D, Yu Y, Holst LP, Luo Y, Fang M, Tamir R, Antonio L, Hu Z, Cupples R, Louis JC, Hu S, Altrock BW and Fox GM. (1996). *Cell*, **85**, 1113–1124.
- Milbrand J, Sauvage FJ, Fahrner TJ, Baloh RH, Leitner ML, Tansey MG, Lampe PA, Heuckeroth RO, Kotzbauer PT, Simburger KS, Golden JP, Davies JP, Vejsada R, Kato AC, Hynes M, Sherman D, Nishimura M, Wang LC, Vandlen R, Moffat B, Klein RD, Poulsen K, Gray C, Garces A, Henderson CE, Phillips HS and Johnson EM. (1998). *Neuron*, **20**, 245–253.
- Mulligan LM, Kwok JB, Healey CS, Elsdon MJ, Eng C, Gardner E, Love DR, Mole SE, Moore JK, Papi L, Ponder MA, Telenius H, Tunnacliffe A and Ponder BAJ. (1993). *Nature*, **363**, 458–460.
- Nakamura T, Ishizaka Y, Nagao M, Hara M and Ishikawa T. (1994). *J. Pathol.*, **172**, 255–260.
- Pachnis V, Mankoo B and Constantini F. (1993). *Development*, **199**, 1005–1017.
- Romeo G, Ronchetto P, Luo Y, Barone V, Seri M, Ceccherini I, Pasini B, Bocciardi R, Lerone M, Kaariainen H and Martucciello G. (1994). *Nature*, **367**, 377–378.
- Santoro M, Carlomagno F, Romano A, Bottaro DP, Dathan NA, Grieco M, Fusco A, Veccio G, Matoskova B, Kraus MH and Di Fiore PP. (1995). *Science*, **267**, 381–383.
- Treanor JJS, Goodman L, De Sauvage F, Stone DM, Poulsen KT, Beck CD, Gray C, Armanini MP, Pollock RA, Hefti F, Phillips HS, Goddard A, Moore MW, Buj Bello A, Davies AM, Asai N, Takahashi M, Vandlen R, Henderson CE and Rosenthal A. (1996). *Nature*, **382**, 80–83.

In summary, the duplication of Cys 634, similar to the substitution in the classical MEN 2A associated mutations, results in an alteration of the number of cysteines which results in spontaneous receptor dimerization. In addition, the different spacing of cysteines may modulate the transformational capacity in various tissues resulting in a different clinical phenotype.

Acknowledgements

We are grateful to M Santoro and A Fusco for making the cDNA of the *RET* short isoform available to us. We thank James Olcese, Ulrike Klein and Natalia Schlambitz for critical reading of the manuscript. The technical assistance of Monika Kistler and the groups of HU Häring (Tübingen) and H Klein (Lübeck) is gratefully acknowledged. This work was supported by Forum Schilddrüse and the Deutsche Forschungsgemeinschaft (Graduiertenkolleg 336).

EXHIBIT B



The Glu632-Leu633 deletion in cysteine rich domain of Ret induces constitutive dimerization and alters the processing of the receptor protein

Italia Bongarzone^{*1}, Elena Vigano¹, Luisella Alberti¹, Piera Mondellini¹, Mauro Uggeri¹, Barbara Pasini², Maria Grazia Borrello¹ and Marco A Pierotti¹

¹Division of Experimental Oncology A, Istituto Nazionale Tumori, 20133 Milan, Italy; ²Scientific Direction, Istituto Nazionale Tumori, 20133 Milan, Italy

Mutations of the *RET* gene, encoding a receptor tyrosine kinase, have been associated with the inherited cancer syndromes MEN 2A and MEN 2B. They have also further been associated with both familial and sporadic medullary thyroid carcinomas. Missense mutations affecting cysteine residues within the extracellular domain of the receptor causes constitutive tyrosine kinase activation through the formation of disulfide-bonded homodimers. We have recently reported that a somatic 6 bp in-frame deletion, originally coding for Glu632-Leu633, potentially activates the *RET* gene. This activation is increased with respect to the frequent MEN 2A-associated missense mutation Cys634Arg. This finding specifically correlated to the clinic behavior of the corresponding tumor, which was characterized by an unusually aggressive progression with both multiple and recurrent metastases. By examining the possibility that this deletion acts in a manner similar to cysteine substitution, we have analysed the molecular mechanism by which this oncogenic activation occurs. Phosphorylated dimers of the deleted Ret receptor were detected in immunoprecipitates separated under non-reducing conditions. Like other Cys point mutations, this 6 bp deletion affecting two amino acid residues between two adjacent Cys, is capable of activating the transforming ability of Ret by promoting receptor dimerization. These results suggest that alteration to cysteine residue position or pairing is capable of inducing ligand independent dimerization. Furthermore, we present data demonstrating that the processing and sorting of the Ret membrane receptor to the cell surface is affected by mutation type.

Keywords: *RET*; oncogenic activation; dimerization

Introduction

Medullary thyroid carcinoma (MTC) is a malignant tumor arising from calcitonin-secreting parafollicular C cells. MTCs may occur sporadically or as a component of the familial cancer syndrome known as multiple endocrine neoplasia type 2 (MEN 2A). Germline mutations within the *RET* proto-oncogene, coding for a receptor tyrosine kinase, can cause the MEN 2A, MEN 2B, and FMTC syndromes. The majority of germline mutations reportedly associated with MEN

2A and FMTC lie within exons 10 and 11 of *RET*. These mutations specifically occur at codons 609, 611, 618, 620 and 634 and universally result in the substitution of a cysteine residue (Mulligan *et al.*, 1993; Donis-Keller *et al.*, 1993). A single point mutation at codon 918, causing the substitution of a methionine (ATG) with a threonine (ACG), is associated with both MEN 2B and sporadic MTC (Eng *et al.*, 1994, 1995b; Carlson *et al.*, 1994; Hofstra *et al.*, 1994; Muragaki *et al.*, 1995; Blaugrund *et al.*, 1994). Approximately 80% of sporadic MTCs are positive for this specific *RET* codon 918 mutation (Eng *et al.*, 1995b, 1996). In contrast, two other relatively rare somatic mutations, at codons 768 (exon 13) and 883 (exon 15), are detected in approximately 10% of sporadic MTCs (Bolino *et al.*, 1995; Eng *et al.*, 1995a; Komminoth *et al.*, 1995). We have recently described two additional interstitial deletions affecting exons 10 and 11 (Romei *et al.*, 1996; Ceccherini *et al.*, 1997). These novel somatic mutations, found in two different patients, were heterozygous in frame deletions of 48 (cod 592–607) and 6 bp (cod 632–633) respectively, yet did not directly involve any cysteine residues. Other similar codon deletions have been observed in sporadic MTCs, namely a 3 bp deletion (cod 633) (Hofstra *et al.*, 1996), and a 27 bp deletion comprising exon 10 (Kalinin and Frilling, 1998). Of interest, a duplication of 12 bp between codon Cys634 and Arg635, resulting in the insertion of four amino acids including a cysteine, has been described as a novel MEN 2A family germline mutation (Hoppner and Ritter, 1997). It has been demonstrated that missense mutations involving any of the cysteine residues 609, 611, 618, 620 and 634, typical of the MEN 2A and FMTC phenotypes yet rarely identified in sporadic MTCs, can promote ligand independent Ret dimerization leading to constitutive enzymatic activation. It is further evident that the substitution of one of these five crucial cysteines leads to both ligand independent dimerization and receptor phosphorylation. We have recently demonstrated in an *in vitro* system that the deletion of the codons 632–633 more effectively activated the *RET* gene with respect to the Cys634Arg missense mutation. This specifically correlated to the clinical severity of the corresponding tumor, characterized by an unusual aggressive progression with both multiple and recurrent metastases (Ceccherini *et al.*, 1997).

In this paper, we addressed the problem of the molecular mechanism underlying this oncogenic activation. We examined the possibility that these deletions act in a similar way to Cys point mutations. Phosphorylated dimers of the deleted Ret receptor

*Correspondence: I Bongarzone

Received 6 November 1998; revised 18 March 1999; accepted 18 March 1999

are detected when immunoprecipitates are separated under non-reducing conditions. Therefore, like Cys point mutations, this deletion affecting two Cys-proximal residues is capable of activating the transforming ability of Ret by promoting ligand-independent receptor dimerization.

Furthermore, we have studied the biogenesis of RetCys634Arg and Retdel1632–633 and found that the latter to be more slowly processed to the cell surface. A large fraction of Retdel632–633 intermediates and a significantly smaller fraction of RetCys634Arg, is retained within the endoplasmic reticulum where it dimerizes and transduces mitogenic signals.

Results

Effect of $\Delta 6$ deletion mutation on dimerization and autophosphorylation of Ret receptor

Unlabeled lysates from Retdel632–633 (Ret $\Delta 6$), RetCys634Arg (Ret2A) and Ret wild-type (Retwt) expressing cell lines were immunoprecipitated with an anti-Ret antibody. Immunocomplexes were tested for kinase activity in the presence of [γ - 32 P]ATP. A significant amount of Ret autophosphorylation was seen in the cell lines expressing the Ret $\Delta 6$ and Ret2A constructs but not in the wild-type (data not shown). This was demonstrated by the labeling of two proteins corresponding to the Ret isoforms of approximately 140 and 160 kDa. Therefore, even in the absence of ligands, both Ret $\Delta 6$ and Ret2A can be phosphorylated.

To examine whether increased kinase activity correlated with increases in covalent dimer formation, we analysed immunoprecipitated Ret $\Delta 6$, Ret2A and

Retwt proteins by SDS–PAGE under both reducing and non-reducing conditions. As shown in Figure 1A, Ret immunoprecipitates separated under non-reducing conditions revealed the presence of Ret dimers in cell lines expressing mutant receptors. Stable dimer formation was not detected in a cell line expressing the wild-type Ret receptor. Furthermore, Ret $\Delta 6$ and Ret2A dimers were shown to be strongly phosphorylated (Figure 1B) by anti-phosphotyrosine immunoblotting. Two high molecular bands corresponding to the dimerized form of the mature (160 kDa) and the partially glycosylated precursor (140 kDa) of the mutated receptor, appeared phosphorylated. Duplicate immunoprecipitates were subjected to immunoblot analysis for Ret after separation under reducing conditions to determine whether receptor dimerization required disulfide bonding (Figure 1C). Results revealed that under reducing conditions, only monomeric forms of Ret were detected. These results provide evidence that the Ret Glu632–Leu633 deletion promotes disulfide-linked dimer formation. Ret proteins were further analysed under non-reducing conditions by 125 I-labeled protein-A binding (data not shown). Densitometric analysis of bands corresponding to dimers or monomers of the two mutant receptors revealed that Ret $\Delta 6$ protein exhibits a greater ability to dimerize in comparison to Ret2A protein. The molar ratio of dimers to monomers for Ret $\Delta 6$ and Ret2A was 0.95 and 0.70 respectively. In contrast, the Retwt control ratio was 0.05. Results from *in vitro* kinase assays, and from both reducing and non-reducing immunoblots demonstrated that the $\Delta 6$ deletion mutation is a gain-of-function mutation. This results in the constitutive activation of Ret, thus allowing for ligand-independent signal transduction.




Figure 1 Dimerization and tyrosine phosphorylation of Ret receptors. Lysates from NIH3T3 cells transfected with the Ret $\Delta 6$ (lanes $\Delta 6$), Ret2A (lanes 2A) and Retwt (lanes wt) constructs were immunoprecipitated (IP) with anti-Ret antiserum (α Ret) and the resulting complexes were separated by SDS–PAGE (6%) under non-reducing (A and B) or reducing conditions (C). Western blot filters were developed (BLOT) with anti-Ret (α Ret, A and C) or anti-phosphotyrosine (α Ptyr, B) antiserum. The positions of the Ret oligomers and monomers are indicated

Effect of 2-mercaptoethanol treatment on Ret mutants transforming activity

Reducing agents have previously been used to interfere with the signaling from activated forms of both the thrombopoietin and NEU receptor (Alexander *et al.*, 1995; Siegel and Muller, 1996). We examined whether the addition of 2-mercaptoethanol (2-ME) could interfere with the ability of the Ret Δ 6 deletion mutant or the Ret2A point mutant to transform NIH3T3 cells. Addition of increasing concentrations of 2-ME (Table 1) failed to affect the transforming activity of the Ret Δ 6 construct. In contrast, Ret2A transforming activity decreased by 84% at the maximal dose (500 μ M) of 2-ME. Toxic effects due to 2-ME at 500 μ M were not evident as demonstrated by the unaffected ability of Ret Δ 6 and HRAS in transforming NIH3T3 cells. This data demonstrates that Ret Δ 6 has a greater resistance to a reducing agent in comparison with Ret2A. This specifically correlates to a stronger disulfide bond formation in the Ret Δ 6 mutant.

Biosynthesis and glycan processing of Ret Δ 6 and Ret2A proteins and effects of 2-mercaptoethanol treatment

It has previously been demonstrated that proteins exposing reactive thiols are often retained within the endoplasmic reticulum (ER) (Isidoro *et al.*, 1996). This protein quality control system has been termed thiol-mediated retention. This mechanism is believed to monitor the oxidation status of one or more cysteines contained in protein sequences. Since Ret Δ 6 and Ret2A mutations affect critical cysteine residues, we studied both the biosynthesis and relative kinetics of dimerization by following the fate of metabolically labeled mutant receptors in NIH3T3 cells. Subconfluent cultures were labeled with 0.35 mCi/ml 35 S-methionine-cysteine for 15 min and subsequently chased for varying times. Under reducing conditions, newly synthesized protein for both mutant receptors appeared in the 140 kDa form immediately after the pulse (Figure 2). The amount of Ret2A p140 protein increased significantly within the first hour of chase and gradually decreased after 2 h. The mature p160 form appeared after 1 h of chase. In contrast, Ret Δ 6 p140 reached a maximum intensity only after 2 h of chase, the same chase time at which the mature p160 form became evident. Since the 140 kDa is the endoglycosidase H sensitive form (data not shown and (Asai *et al.*, 1995)) whilst the 160 kDa is resistant, we deduced the longer persistence of the Ret Δ 6 140 kDa precursor. This indicated that Ret Δ 6 has a less efficient ER processing compared to the Ret2A precursor. The amount of the mature 160 kDa Ret2A protein was similar to that of the precursor after 2 h of chase, with a maximum reached after 4 h. In contrast, the amount of the 160 kDa Ret Δ 6 protein increased slowly, yet always remained less than the 140 kDa form. To explore the mechanism of dimerization, features of the labeled Ret Δ 6 and Ret2A proteins were also followed as a function of time by non-reducing SDS-PAGE (Figure 2). Both the precursor and mature forms of Ret Δ 6 and Ret2A were capable of dimerization, each with individual specific kinetics (Figure 2). The formation of Ret Δ 6 homodimers in

Table 1 2-mercaptoethanol effect on the transforming activity of Ret mutants

Transfected plasmid (125 ng of DNA)	Transformed foci/plate at different concentration of 2-ME (μ M)			
	0	30	125	500
pRCRET Δ 6	179 \pm 11	178 \pm 7	185 \pm 26	180 \pm 19
pRCRET2A	87 \pm 6	87 \pm 9	89 \pm 11	14 \pm 7
pRCRETwt	—	—	—	—
pHRAS	152 \pm 18	172 \pm 11	179 \pm 7	160 \pm 4

Focus assays were performed with 125 ng of plasmid per plate in the presence of increasing concentrations of 2-ME. Foci were scored over six plates for each concentration after 3 weeks. The HRAS oncogene mutated at its 12th codon was used as a control. Values listed represent the average \pm s.d. of the number of foci counted on six plates




Figure 2 Pulse-chase labeling of Ret Δ 6 and Ret2A proteins in reducing and non-reducing conditions. Autoradiograms of metabolically labeled radioactive Ret Δ 6 and Ret2A proteins derived from transfectants pulsed for 15 min and harvested immediately (lanes marked 0) or after the indicated chase time (lanes marked 1–6). Cell extracts were immunoprecipitated with anti-Ret antiserum and the complexes were separated by SDS-PAGE under either reducing or non-reducing conditions. Gels were exposed to X-ray film for 4–5 days. Numbers are molecular weights in kDa

both the p140 and p160 forms was observed from 2 h. In contrast, after 1 h of chase in the Ret2A mutant, a consistent amount of p140 homodimers and only a small fraction of p160 homodimers were observed. Furthermore, p160 homodimers were observed almost exclusively after longer chase times. Analysis of phosphorylation levels in all Ret Δ 6 and Ret2A proteins indicated that most of the precursors were significantly phosphorylated, irrespective of whether in the monomeric or dimeric form (Figures 3B and 1B and data not shown). This is due to the fact that phosphorylation associated to dimer forms is intermolecular. Therefore, SDS-PAGE under reducing conditions produces phosphorylated monomers.

Co-immunoprecipitation of SHC phosphorylated proteins with both the precursor and mature forms of the Ret Δ 6 and Ret2A receptors

Ret Δ 6 proteins are characterized by slower maturation kinetics yet display stronger oncogenic activity in comparison to Ret2A proteins. With the aim to explain this fact, we speculated that both the p140 precursor and p160 mature forms of Ret, when




Figure 3 Coimmunoprecipitation of Shc phosphorylation proteins with both the precursor and the mature forms of Ret Δ 6 and Ret2A receptors. Lysates from NIH3T3 cells (lane NIH) or from NIH3T3 cell lines expressing the Ret Δ 6 (lane Δ 6) and Ret2A (lane 2A) proteins were immunoprecipitated (IP) with anti-Shc (α Shc, A and B) or anti-Ret (α Ret, C and D) antiserum. The resulting complexes were electrophoresed on a 7.5% SDS-PAGE under reducing conditions. Proteins were transferred to nitrocellulose and subjected to immunoblot analysis (BLOT) with anti-phosphotyrosine (α Ptyr) or anti-Ret (α Ret) antiserum. Ret Δ 6 and Ret2A expressing cell lines were serum starved for 48 h. Numbers are molecular weights in kDa and correspond to two of the three different isoforms of Shc (p52 and p46) and to Ret (p160 and p140), respectively

phosphorylated and dimerized, are able to transduce mitogenic activity in spite of their different localization; the endoplasmic reticulum or surface membrane respectively. To address this, we investigated whether the precursor forms of Ret Δ 6 and Ret2A were able to bind the Shc adapter protein. This protein is localized to the membrane of the rough endoplasmic reticulum and has been previously demonstrated to bind Ret/PTCs, Ret2A and Ret2B (a mutant Ret receptor carrying an M918T mutation) oncoproteins (Lotti *et al.*, 1996). Cell lysates from Ret Δ 6 and Ret2A transfected NIH3T3 cells were immunoprecipitated with anti-Shc or anti-Ret antibodies and subsequently blotted with anti-Ptyr antibodies (Figure 3). Both the precursor and mature forms of Ret were co-immunoprecipitated with Shc phosphorylated proteins. The band density of both Ret Δ 6 and Ret2A co-immunoprecipitated or not with Shc protein and probed with anti-Ptyr were checked by densitometric analysis. The observed affinity of Shc for the precursor and mature forms of both the mutated receptors is similar. Considering the sum of the relative amounts, it was calculated that 40% of Ret Δ 6 and 20% of Ret2A were co-immunoprecipitated with Shc phosphorylated proteins. It is thus possible to conclude from this data that the mitogenic signal transduced by Shc does not depend on maturation or localization of the mutated receptor, but possibly influenced by the specific type of activating mutation.

Discussion

We have demonstrated that deletion of the residues Glu632-Leu633 of Ret induces stable receptor dimer formation in the absence of ligand. This suggests that constitutive dimerization and autophosphorylation of the receptor protein underlies the potent transforming activity of this Ret mutant. MEN 2A mutations destroy Cys residues, thus indicating that their

mechanism of action involves the creation of ligand-independent dimers most likely through aberrant intermolecular disulfide-bond formation. Whilst not directly involving Cys residues, the Δ 6 mutation also induces such dimerization. This deletion mutation may indirectly affect neighboring Cys residues by disrupting the normal and most likely intramolecular cysteine pairing, which occurs in the non-activated receptor. This thus leaves the possibility that at least one unpaired cysteine residue may participate in an intermolecular binding event with another altered receptor, resulting in a disulfide-bonded receptor dimer leading to aberrant receptor activation. In fact, a similar mechanism has been proposed to account for the oncogenic activation in Neu deletion mutants. Deletion of conserved cysteine residues in the juxtatransmembrane region of Neu resulted in elevated transforming activities compared to that of the C647S point mutation form (Siege and Muller, 1996). One potential explanation for this observation is that the spatial arrangement between the remaining cysteine residues may play a crucial role in mediating receptor dimerization. For example, certain deletions may result in a free cysteine being exposed at the surface of the receptor, thus making it more accessible to intermolecular disulfide bond formation.

Interestingly, the transforming activity of Ret2A receptor is decreased following the addition of a reducing agent. In contrast, the transforming activity of the Ret Δ 6 deletion mutant is not significantly impaired at the same non toxic concentration of 2-ME. This may be explained by the formation of more stable Ret Δ 6 disulfide-linked dimers, thus generating a more efficient covalent dimerization. Alternatively, it is possible that the loss of residues near cysteine 634 results in an unfolding of the cysteine rich domain that may lead to a less compact and more flexible conformation. Such an unfolding could increase the probability of receptor interaction on the plasma membrane, leading to the tyrosine kinase activation

of a greater number of mutant receptors compared to the Ret2A type receptor. Finally, the lower sensitivity of the $\Delta 6$ mutant to reducing agents could also be explained by its particular molecular equilibrium shift from monomeric to dimeric/oligomeric forms.

Taken together, these observations suggest that both the number and spatial arrangement of cysteine residues may induce a ligand independent dimerization.

More generally, it is possible that the loss, gain or mutation of residues within the cysteine-rich region of Ret has the capacity to induce stabilization of the dimeric conformation through intermolecular disulfide bonds.

Biosynthesis studies of Ret $\Delta 6$ and Ret2A proteins have shown a slower rate of maturation for Ret $\Delta 6$ protein due to precursor retention in the endoplasmic reticulum. However, both Ret $\Delta 6$ and Ret2A show intracellular retention in comparison with the Retwt protein, which reaches the membrane within a 30 min chase time (data not shown). This is expected since, as a rule, only proteins that have attained a proper three-dimensional structure are secreted or expressed at the cell surface. Since Ret $\Delta 6$ and Ret2A are mutated near or within Cys, their folding and assembly could potentially be delayed by intracellular retention for a longer time with respect to Retwt.

Accumulation of misfolded immature Ret2A and Ret $\Delta 6$ proteins in the intracellular vesicular compartment and its homodimerization due to aberrant intermolecular disulfide bonds have important consequences on receptor functions in the downstream mitogenic program. In fact, a very large portion of Ret $\Delta 6$ and Ret2A precursors strongly bind phosphorylated Shc proteins, thus indicating their ability to induce this mitogenic signal pathway. This finding may also aid in explaining why the Ret $\Delta 6$ protein is more oncogenic, even when less efficiently processed than Ret2A mutant; an obvious paradox. The possibility that intracellular Ret precursors still trigger a mitogenic activity is extremely important also in the context of the effects of cysteine mutation in positions 609, 618 and 620. There is evidence suggesting that these cysteine mutations induce constitutive catalytic activity due to aberrant disulfide homodimerization of Ret, but in the same time are responsible for a decrease in the amount of mature RET protein expressed at the cell surface (Chappuis-Flament *et al.*, 1998). Since 20–30% of families with mutations C618R or C620R (Mulligan *et al.*, 1994) present a combination of HSRC and MEN 2A, it has been hypothesized that these cysteine mutations exert a dual impact on RET. In particular, these mutations can be activating or deactivating depending on the tissue in which RET is expressed. Thus, RET carrying either the C618R or C620R mutation, due to constitutive dimerization, exhibits a transforming potential responsible for MEN2A and in the same time, due to scarce localization at the surface membrane, an inability to promote a survival of enteric neurons. Our data well supports this model, indicating that not exclusively the membrane can trigger mitogenic signals. This reinforces the concept that an efficient membrane localization of Ret receptors is very important for triggering of neuronal differentiation signals.

Materials and methods

Cell culture and transfection

Mouse NIH3T3 cells were grown in Dulbecco's modified Eagle's medium (DMEM) containing 10% calf serum. Transfection experiments were performed in NIH3T3 cells by calcium phosphate precipitation as previously described (Bongarzone *et al.*, 1993). Plasmids carrying Retwt, Ret $\Delta 6$ and Ret2A inserts have been described in detail previously (Ceccherini *et al.*, 1997). The HRAS oncogene was mutated at its 12th codon by a G→T transversion. Transformation foci were selected in DMEM plus 5% calf serum. The reducing agent 2-ME was added when cells had formed a monolayer. This monolayer was maintained for 14 days in DMEM supplemented with 5% bovine calf serum.

Immunoprecipitation and Western blot analysis

Protein samples were prepared as previously described (Borrello *et al.*, 1996) and immunoprecipitated with the specified antisera: affinity purified anti-Ret polyclonal antiserum (Borrello *et al.*, 1996) and rabbit polyclonal anti-Shc antiserum (Upstate Biotechnology Incorporated). Immunoprecipitates were resolved by electrophoresis on 7.5 or 8.5% SDS polyacrylamide gels (PAGE). Proteins were transferred to nitrocellulose filters, blocked with 5% bovine serum albumin (BSA) or 0.5% gelatin in Tris-buffered saline (TBS) pH 7.6 and immunoblotted with the same anti-Ret antiserum described above or with the monoclonal antiserum anti-phosphotyrosine (anti-Ptyr, Upstate Biotechnology Incorporated). Immunoreactive bands were visualized using horseradish peroxidase conjugated anti-rabbit or anti-mouse antisera and ECL detection reagents (Amersham) or using 125 I-labeled protein A (Amersham) followed by autoradiography. When 125 I-labeled protein A was used, filters were exposed to storage phosphor screen films (Molecular Dynamics) in order to quantify the counts per minute (c.p.m.) associated with the Ret-specific bands reacting with anti-Ret or anti-Shc with a Phosphorimager. For Western blotting under non-reducing conditions, 2-mercaptoethanol was excluded.

Metabolic labeling and immunoprecipitation

Prior to labeling, subconfluent cells were incubated with methionine-free DMEM for 1 h. Metabolically labeling was performed with 370 μ Ci/mL 35 S-Methionine-Cysteine (DuPont/NEN) for 15 min. The labeling medium was then removed and replaced with completed DMEM medium for the duration of the chase period. Further processing, extraction, and immunoprecipitation of labeled Ret was carried out as above with the following modifications: after incubation, bound immune complexes (Protein A/G-Plus-Agarose beads, Santa Cruz Biotechnology) were washed extensively (seven times) with lysis buffer (0.15 M NaCl/0.05 M Tris-HCl, pH 7.2/1% Triton X-100/1% sodium deoxycholate/0.1% SDS), either supplemented with or without 1% Triton and 1 M NaCl and on a shaking platform, eluted, and subjected to SDS-PAGE fractionation and autoradiography. Densities of radioactive bands on X-ray films were estimated using a Phosphorimager.

Acknowledgements

This work was partially supported by the Associazione Italiana per la Ricerca sul Cancro (AIRC), Fondazione Italiana per la Ricerca sul Cancro (FIRC), CNR (Biotechnologie) No. 97.01258.PF49, Project BIOMED2 No. BMH4-CT97-2157 and Istituto Superiore di Sanita'. The authors thank Dr. R. Sitia for critical comments and suggestions on the manuscript. We also thank Cristina Mazzadi for secretarial help.

References

- Alexander WS, Metcalf D and Dunn AR. (1995). *EMBO J.*, **14**, 5569–5578.
- Asai N, Iwashita T, Matsuyama M and Takahashi M. (1995). *Mol. Cell. Biol.*, **15**, 1613–1619.
- Blaugrund JE, Johns MM, Eby JY, Ball DW, Baylin SB, Hruban RH and Sidransky D. (1994). *Hum. Mol. Genet.*, **3**, 1895–1897.
- Bolino A, Schuffenecker I, Luo Y, Seri M, Silengo M, Tocco T, Chabrier G, Houdent C, Murat A, Schlumberger M, Towniaire J, Senoir GM and Omeo G. (1995). *Oncogene*, **10**, 2415–2419.
- Bongarzone I, Monzini N, Borrello MG, Carcano C, Ferraresi G, Arighi E, Mondellini P, Della Porta G and Pierotti MA. (1993). *Mol. Cell. Biol.*, **13**, 358–366.
- Borrello MG, Alberti L, Arighi E, Bongarzone I, Battistini C, Bardelli A, Pasini B, Piutti C, Rizzetti MG, Mondellini P, Radice MT and Pierotti MA. (1996). *Mol. Cell. Biol.*, **16**, 2151–2163.
- Carlson KM, Dou S, Chi D, Scavarda NJ, Toshima K, Jackson CE, Wells Jr SA, Goodfellow P and Donis-Keller H. (1994). *Proc. Natl. Acad. Sci. USA*, **91**, 1579–1583.
- Ceccherini I, Pasini B, Pacini F, Gullo M, Bongarzone I, Romei C, Santamaria G, Matera I, Mondellini P, Scopsi L, Pinchera A, Pierotti MA and Romeo G. (1997). *Oncogene*, **14**, 2609–2612.
- Chappuis-Flament S, Pasini A, De Vita G, Segouffin-Cariou C, Fusco A, Attie T, Lenoir GM, Santoro M and Billaud M (1998). *Oncogene*, **17**, 2851–2861.
- Donis-Keller H, Dou S, Chi D, Carlson KM, Toshima K, Lairmore TC, Howe JR, Moley JF, Goodfellow P and Wells Jr SA. (1993). *Human Mol. Genet.*, **2**, 851–856.
- Eng C, Mulligan LM, Healey CS, Houghton C, Frilling A, Raue F, Thomas GA and Ponder BA. (1996). *Cancer Res.*, **56**, 2167–2170.
- Eng C, Smith DP, Mulligan LM, Healey CS, Zvelebil MJ, Stonehouse TJ, Ponder MA, Jackson CE, Waterfield MD and Ponder BAJ. (1995a). *Oncogene*, **10**, 509–513.
- Eng C, Mulligan LM, Smith DP, Healey CS, Frilling A, Raue F, Neumann HPH, Pfragner R, Behmel A, Lorenzo MJ, Stonehouse TJ, Ponder MA and Ponder BAJ. (1995b). *Genes Chrom. Cancer*, **12**, 209–212.
- Eng C, Smith DP, Mulligan LM, Nagai MA, Healey CS, Ponder MA, Gardner E, Scheumann GFW, Jackson CE, Tunnacliffe A and Ponder BAJ. (1994). *Human Mol. Genet.*, **3**, 237–241.
- Hofstra RM, Landsvater RM, Ceccherini I, Stulp RP, Stelwagen T, Luo Y, Pasini B, Hoppener JWM, Ploos van Hamstel HK, Romeo G, Lips CJM and Buys CHCM. (1994). *Nature*, **367**, 375–376.
- Hofstra RM, Stelwagen T, Stulp RP, De Jong D, Hulsbeek M, Kamsteeg EJ, van den Berg A, Landsvater RM, Vermey A, Molenaar WM, Lips CJ and Buys CH. (1996). *J. Clin. Endocrin. Metabol.*, **81**, 2881–2884.
- Hoppner W and Ritter MM. (1997). *Human Mol. Genet.*, **6**, 587–590.
- Isidoro C, Maggioni C, Demoz M, Pizzigalli A, Fra AM and Sitia R. (1996). *J. Biol. Chem.*, **271**, 26138–26142.
- Kalinin V and Frilling A. (1998). *J. Mol. Med.*, (in press).
- Komminoth P, Kunz EK, Matias-Guiu X, Hiort O, Christiansen G, Colomer A, Roth J and Heitz PU. (1995). *Cancer*, **76**, 479–489.
- Lotti LV, Lanfrancone L, Igliaccio E, Ompetta C, Elicci G, Alcini AE, Alini B, Elicci PG and Orrisi MR. (1996). *Mol. Cell. Biol.*, **16**, 1946–1954.
- Mulligan LM, Eng C, Attie T, Lyonnet S, Marsh DJ, Hyland VJ, Robinson BG, Frilling A, Verellen-Dumoulin C and Safar A. (1994). *Human Molec. Genet.*, **3**, 2163–2167.
- Mulligan LM, Kwok JBJ, Healey CS, Elsdon MJ, Eng C, Gardner E, Love DR, Mole SE, Moore JK, Papl L, Ponder MA, Telenius H, Tunnacliffe A and Ponder BAJ. (1993). *Nature*, **363**, 458–460.
- Muragaki Y, Timothy N, Leight S, Hempstead BL, Chao MV, Trojanowski JQ and Lee VM. (1995). *J. Comparat. Neurol.*, **356**, 387–397.
- Romei C, Elisei R, Pinchera A, Ceccherini I, Molinaro E, Mancusi F, Martino E, Romeo G and Pacini F. (1996). *J. Clin. Endocrin. Metabol.*, **81**, 1619–1622.
- Siegel PM and Muller WJ. (1996). *Proc. Natl. Acad. Sci. USA*, **93**, 8878–8883.

EXHIBIT C

Ligand Stimulation of a Ret Chimeric Receptor Carrying the Activating Mutation Responsible for the Multiple Endocrine Neoplasia Type 2B*

(Received for publication, July 19, 1996, and in revised form, August 28, 1996)

Claudia Rizzo†, Daniela Califano§, G. Luca Colucci-D'Amato, Gabriella De Vita, Amelia D'Alessio§, Nina A. Dathan¶, Alfredo Fusco||, Carmen Monaco‡§, Giovanni Santelli§, Giancarlo Vecchio, Massimo Santoro, and Vittorio de Franciscis**

From the Centro di Endocrinologia ed Oncologia Sperimentale del CNR, c/o Dipartimento di Biologia e Patologia Cellulare e Molecolare, Università di Napoli "Federico II," via S. Pansini 5, 80131 Napoli, Italy, §Oncologia Sperimentale "E," Istituto Nazionale Tumori, Fondazione "G. Pascale," via M. Semmola, 80131 Napoli, Italy, and ||Dipartimento di Medicina Sperimentale e Clinica, Università degli Studi di Reggio Calabria, via T. Campanella 5, 88100 Catanzaro, Italy

Inherited activating mutations of Ret, a receptor tyrosine kinase, predispose to multiple endocrine neoplasia (MEN) types 2A and 2B and familial medullary thyroid carcinoma. To investigate the effects induced by acute stimulation of Ret, we transfected both PC12 and NIH 3T3 cells with a molecular construct in which the ligand-binding domain of the epidermal growth factor receptor was fused to the catalytic domain of Ret. Acute stimulation of the chimeric receptor induced PC12 cells to express a neuronal-like phenotype. Moreover, we introduced the dominant mutation, responsible for the multiple endocrine neoplasia type 2B, in the catalytic domain of the Ret chimera. Expression of the mutant chimera, in the absence of ligand stimulation, induces the PC12 cells to acquire a flat morphology with short neuritic processes and transforms the NIH 3T3 cells. Stimulation of the mutant chimera with epidermal growth factor causes a drastic overgrowth of long neuritic processes, with the induction of the *suc1*-associated protein tyrosine phosphorylation in PC12 cells and higher transforming efficiency in NIH 3T3 cells. These data indicate that the gain-of-function MEN2B mutation does not abrogate ligand responsiveness of Ret and suggest that the presence of Ret ligand could play a role in the pathogenesis of the MEN2B syndrome.

Specific mutations of the *ret* gene, a receptor tyrosine kinase (1), are responsible for the inheritance of multiple endocrine neoplasia (MEN)¹ type 2A and 2B and familial medullary thyroid carcinoma syndromes (2). MEN2A and MEN2B are distinct hereditary neoplastic syndromes both characterized by the presence of medullary thyroid carcinomas and pheochro-

mocytomas. MEN2A is also characterized by hyperplasia of parathyroid cells, whereas MEN2B is a more severe disease, being associated with skeletal abnormalities, ganglioneuromas of the intestinal tract, and mucosal neuromas, and it is characterized by an earlier age of tumor onset (3). Mutations in cysteine residues of the extracellular domain are the causative genetic event of familial medullary thyroid carcinoma and MEN2A syndromes (4, 5). A single point mutation, which results in a Thr for Met substitution at codon 918 within the Ret catalytic domain, is responsible for the MEN2B syndrome (6, 7). These mutations convert Ret into a dominant transforming gene (*ret*MEN2A and *ret*MEN2B alleles) and cause constitutive activation of its intrinsic tyrosine kinase activity (8, 9).

*ret*MEN2A and *ret*MEN2B differ in their mechanisms of activation. In the case of *ret*MEN2A, activation likely results from constitutive receptor dimerization, whereas *ret*MEN2B proteins do not constitutively dimerize and display altered substrate specificity (2, 9, 10). It is presently unknown whether the Ret harboring the MEN2B mutation is fully activated by an intramolecular mechanism. Indeed, if *ret*MEN2B is still sensitive to ligand stimulation, the contribution of active Ret to the resulting phenotype, in the affected tissues, may be in part attributed to the presence of available Ret ligand in the extracellular environment.

We thus investigated the biological effects induced by Ret stimulation in the rat pheochromocytoma cell line, PC12, because of the sensitivity of this system, which retains the ability to differentiate *in vitro* and also allows discrimination among stimuli from different extracellular signals (11, 12). Indeed, this cell line has been particularly suitable for studying the molecular mechanisms by which *ret* alleles contribute to the development of neuroendocrine cancer syndromes (13–16). We have recently shown that chronic expression of active Ret oncoproteins induces the PC12 cells to differentiate toward a neuronal-like phenotype. Yet, we have shown that Ret-induced differentiation is not complete, because the expression of neuronal genes is dissociated from the inhibition of cell proliferation (16).

Because one of the biological mechanisms underlying the choice between differentiation and proliferation in PC12 cells is determined by the extent and duration of the signaling (12), we decided to investigate whether acute stimulation of Ret causes differentiation of the PC12 cells. In addition, we addressed the question of whether constitutive activation, induced by the MEN2B mutation, fully activates the Ret biochemical activity, thus abrogating responsiveness to ligand stimulation.

A potential physiological ligand for Ret has recently been

* This work was supported in part by grants from the Associazione Italiana per la Ricerca sul Cancro (AIRC) and by the targeted project PF-CNR, ACRO (sottoprogetto 7). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

† Recipient of an Applicazione Clinica della Ricerca Oncologica fellowship.

¶ Recipient of a European Communities fellowship in the Biotechnology program.

** To whom correspondence should be addressed: Centro di Endocrinologia ed Oncologia Sperimentale del CNR, Facoltà di Medicina e Chirurgia, via S. Pansini 5, 80131 Napoli, Italy. Fax: 39-81-7463037; E-mail: defranciscis@cids.unina.it.

¹ The abbreviations used are: MEN, multiple endocrine neoplasia; GDNF, glial cell line-derived neurotrophic factor; EGF, epidermal growth factor; EGFR, EGF receptor; NGF, nerve growth factor.

identified as the glial cell line-derived neurotrophic factor, GDNF (17–20). Ret association to GDNF and its subsequent tyrosine phosphorylation is mediated by the presence in the same cell surface complex of the GDNF receptor- α , a glycosyl-phosphatidylinositol-anchored protein. This protein is expressed in GDNF-responsive tissues and in cultured embryonic neurons, whereas in the cell lines examined, including PC12 and NIH 3T3 cells, complete stimulation of Ret by GDNF depends on the exogenous addition of GDNF receptor- α (19, 20). Thus, to perform experiments in PC12 cells, we utilized a chimeric receptor, EGFR/*ret*, which consists of the transmembrane and ligand-binding domains of the epidermal growth factor receptor (EGFR) fused to the catalytic domain of Ret. Such a chimeric construct has already been shown to be a useful tool in characterizing the Ret-specific transducing signaling (21, 22). In addition, we also utilized a mutant chimera, consisting of the EGFR/*ret* construct in which we introduced a single point mutation, which converts the Met codon 918 (1) into Thr (EGFR/*ret*^{Thr-918} chimera).

The results reported here indicate that the MEN2B mutation does not cause full activation of Ret, since it retains the ability to be further stimulated by an extracellular ligand. This stimulation resulted in an increased autophosphorylation of the receptor, pronounced neurite outgrowth, and tyrosine phosphorylation of Snt. Thus, our results suggest that the MEN2B disease phenotype could be influenced by the tissue distribution of a Ret ligand.

EXPERIMENTAL PROCEDURES

Cell Culture and Transfection Experiments—PC12 cells were grown in RPMI 1640 (Life Technologies, Inc.) supplemented with 10% horse serum and 5% fetal calf serum (13). NIH 3T3 cells were grown in DMEM supplemented with 10% fetal calf serum (21). Transfection experiments were performed with 10 μ g of plasmid DNA using either the Lipofectin reagent (Life Technologies, Inc.) for PC12 cells (13) or the calcium phosphate precipitation method for NIH 3T3 cells. The transfected cells were selected in *gpt* selection medium for 3 weeks, and individual cell colonies were isolated and expanded. Epidermal growth factor (Upstate Biotechnologies, Inc. (UBI)) or 2.5 S nerve growth factor (UBI) (100 ng/ml) were added to the culture medium as indicated.

Immunoprecipitation and Immunoblotting—Between 10⁶ and 10⁷ cells were washed twice in ice-cold Tris-buffered saline (20 mM Tris-HCl, pH 8.0, 150 mM NaCl) and then lysed in a buffer containing 50 mM HEPES, pH 7.5, 1% (v/v) Triton X-100, 50 mM NaCl, 5 mM EGTA, 50 mM NaF, 20 mM sodium pyrophosphate, 1 mM sodium orthovanadate, 2 mM phenylmethylsulfonyl fluoride, 0.2 μ g/ml each of aprotinin and leupeptin, and 4 mM diisopropylfluorophosphate and clarified by centrifugation at 10,000 $\times g$ for 15 min, as previously reported (21). Protein concentrations were estimated by a modified Bradford assay (Bio-Rad). Equal amounts of protein were incubated with rabbit anti-Ret antibody, as indicated, for 1 h at 4 °C and subsequently incubated with protein A-Sepharose CL4-B (Pharmacia) for 1 h at 4 °C. Immunoprecipitates were washed three times with the above mentioned lysis buffer and boiled in Laemmli buffer for 5 min before electrophoresis. Immunoprecipitates were subjected to SDS-PAGE (7.5% polyacrylamide) under reducing conditions and transferred to polyvinylidene difluoride (Millipore Corp.). Immunoblotting was carried out using either anti-Ret antibodies or anti-phosphotyrosine monoclonal antibodies (UBI, G410), and the reaction was detected with peroxidase-conjugated secondary antibodies and Amersham ECL system. The polyclonal antibody (anti-Ret) was generated against a fusion protein in which the tyrosine kinase domain of human Ret is fused to the bacterial glutathione S-transferase (21). The Snt protein was isolated from cell lysates using p13^{suc1}-agarose (Oncogene Science) as described (23).

Northern Blot Analysis—RNA was prepared from cultured cells by a modification of the guanidine thiocyanate method (24). 20 μ g of total RNA were size-fractionated on a denaturing formaldehyde agarose gel and blotted onto nylon filters (Hybond-N, Amersham Corp.). To obtain the *krox24* probe, 60-mer oligonucleotides were synthesized according to the published sequence and subsequently ³²P-labeled using the Klenow fragment of the *Escherichia coli* DNA polymerase and a 3'-terminal specific 9-mer. The *vgf* probe was excised from the pV2-2 plasmid (25). ³²P labeling of the *vgf* and ribosomal 18 S probes was performed

with the random oligonucleotide primer kit (Amersham). Hybridization and washing were carried out under stringent conditions: 0.1 \times SSC, 0.1% SDS, 60 °C. Autoradiography was performed using Kodak X-AR films at -70 °C for 1–7 days with intensifying screens.

RESULTS

Acute Stimulation of Ret Induces Neurite Outgrowth in PC12 Cells—PC12 cells, stably transfected with the chimera (EGFR/*ret*) or with the vector alone (LTR-3) (Fig. 1A), were selected for resistance to mycophenolic acid. A mass population and individual clones were then isolated and analyzed. All of the EGFR/*ret*- and vector-transfected populations appeared morphologically undifferentiated, displaying a small size and round shaped morphology (Fig. 1B and data not shown).

Stimulation of the EGFR/*ret* chimera with epidermal growth factor (EGF) (100 ng/ml) induced the PC12 cells to change, within 24 h, from a round shaped to a neuron-like morphology with long neurite processes that strikingly resembled that induced by the nerve growth factor (NGF) on the parental cells (Fig. 1B). On the other hand, EGF stimulation (up to 300 ng/ml) of the parental cells and of the vector-transfected cells had little or no effect on cell morphology even after 72 h of continuous treatment (Fig. 1B and data not shown).

The expression and the functional integrity of the EGFR/*ret* chimeric receptor were tested by immunoprecipitation of Ret products, followed by blotting either with anti-Ret or with anti-phosphotyrosine antibodies (Fig. 1C). A single protein, of 140–150-kDa apparent molecular mass, corresponding to the EGFR/*ret* product (21), was observed in the PC12-EGF/*ret* cells but not in the parental cells. In the absence of ligand stimulation, the EGFR/*ret* receptor displayed some constitutive levels of phosphorylation; however, stimulation with EGF (100 ng/ml) caused a dramatic increase (more than 20-fold) in tyrosine phosphorylation of the receptor (Fig. 1C).

Ret Induces Differentiation in PC12 Cells—NGF-induced differentiation of the PC12 cells is characterized by the expression of a complex pattern of genes, including immediate early genes (*fos*, *krox24*) or *delayed and late genes* (*vgf*, SCG10, peripherin), the expression of the latter genes being, at least partially, dependent on protein synthesis (26). We decided to investigate whether the Ret-induced neurite outgrowth was associated with the expression of a similar pattern of genes. In Fig. 2, we show that stimulation of the chimera induced the expression of *krox24* and *vgf* (27, 28) at levels similar to those induced by NGF. On the other hand, EGF stimulation was unable to induce any significant *vgf* expression, and it induced only low levels of *krox24* mRNA.

Because EGF is able, on its own, to induce an early gene response, which partially superimposes that induced by NGF, even if to a lower extent, we determined whether stimulation of the chimera could induce tyrosine phosphorylation of a specific target of neurotrophic factor activity in the neuronal cells, Snt (Fig. 3). Tyrosine phosphorylation of Snt has been reported as a qualitative event that discriminates between proliferation signals, induced by serum or EGF, and differentiation signals, such as that induced by NGF, even if its biological function is still poorly understood (23, 29). EGF stimulation of the EGFR/*ret* chimera resulted in tyrosine phosphorylation of Snt at levels similar to those observed following stimulation with NGF (Fig. 3, compare lane 5 to lanes 3 and 6). On the other hand, EGF had no effect on the parental PC12 cells (Fig. 3, lane 2).

Ligand Stimulation Increases the retMEN2B Activity in PC12 and NIH 3T3 Cells—A single point mutation in the catalytic domain of Ret, which is associated with the MEN2B syndrome, causes constitutive tyrosine kinase activation. This mutation enables Ret to transform the NIH 3T3 cell line and to cause incomplete differentiation of the PC12 cells (9, 16). To




FIG. 1. Panel A, schematic representation of the EGFR/*ret* construct. The EGFR/*ret* chimera encompasses the extracellular and transmembrane domains of the EGFR, and the intracellular domain of Ret (21). Met-918 is also indicated. Panels B and C, effects of ligand-dependent stimulation of the EGFR/*ret* chimera on PC12 cell morphology and phosphorylation of the receptor. Panel B, phase contrast micrograph of parental PC12 cells (upper parts) or EGFR/*ret* (lower parts) transfectants. Cells were grown for 72 h in the presence of either NGF (100 ng/ml) or EGF (100 ng/ml) or left untreated as indicated. EGF induces neurite outgrowth in EGFR/*ret* transfectants, which are similar to those observed in the parental cells treated with NGF. Panel C, total cellular proteins were immunoprecipitated (Ipt) with the anti-Ret polyclonal antibody (α ret) and then divided into two aliquots and analyzed by immunoblotting (blot) with the anti-Ret or anti-Tyr(P) (α pTyr) antibody, as indicated. Control mock-transfected PC12 cells (C) or EGFR/*ret* transfectant (E/R) were either untreated (–) or treated with 100 ng/ml EGF for 5 min at 37 °C (+). Molecular mass markers are indicated in kilodaltons. The position of the EGFR/*ret* chimera is also indicated.

address the question of whether the *ret*MEN2B is further inducible, we introduced the Met-918 to Thr substitution in the EGFR/*ret* construct (thus obtaining the EGFR/*ret*^{Thr-918}). We first evaluated its effects in NIH 3T3 cells in a focus formation assay. Consistent with the “gain of function” effects of the MEN2B mutation (9), EGFR/*ret*^{Thr-918} transformed also in the absence of EGF (10² focus-forming units/pmol). EGF stimulation further increased the transforming activity of the EGFR/*ret*^{Thr-918} construct (Table I), indicating that also in the presence of a MEN2B mutation, Ret retained responsiveness to ligand triggering.

PC12 cells were then transfected with the EGFR/*ret*^{Thr-918} construct, and both a mass population and several independent clones were marker-selected. The morphology of PC12-EGFR/*ret*^{Thr-918} cells was indistinguishable from that previously reported in the case of PC12 cells expressing a *ret*MEN2B allele (16). Indeed, PC12-EGFR/*ret*^{Thr-918} cells were flat and showed the growth of short neurites (Fig. 4). However, they were still responsive to ligand triggering. Twenty-four hours of EGF treatment induced the PC12-EGFR/*ret*^{Thr-918} cells to shift to-

ward a more differentiated neuronal phenotype that was, however, clearly different from that induced by NGF on parental cells. As shown in Fig. 4, although EGF induced a pronounced neuritic outgrowth, PC12-EGFR/*ret*^{Thr-918} cells still retained a flat shaped cell body that contrasted with the round shape and the high refractility characterizing PC12 cells treated with NGF.

These biological effects were explained by the retained responsiveness of the tyrosine kinase activity of the EGFR/*ret*^{Thr-918} construct to ligand stimulation. Consistent with the reported constitutive activation of the tyrosine kinase function of Ret caused by the MEN2B mutation (9), the EGFR/*ret*^{Thr-918} protein product showed constitutive levels of tyrosine phosphorylation, both in NIH 3T3 and PC12 cells, which were higher than those of the wild type EGFR/*ret* chimera (Fig. 5A and data not shown). However, EGF stimulation caused a sharp increase in the phosphorylation of the receptor in both cell lines. On the other hand, phosphorylation of Snt seems to correlate with the levels of tyrosine phosphorylation of the receptor. In fact, Snt was barely phosphorylated in cells transfected with EGFR/




FIG. 2. Gene expression induced by EGFR/ret in PC12 cells. Northern blot analysis is shown of total cellular RNA (20 μ g) extracted either from PC12 cells (lanes 1–5) or from PC12 cells transfected with EGFR/ret (lanes 6–8), grown in the presence of NGF (100 ng/ml) or EGF (100 ng/ml) as indicated. The filters were hybridized with either a *krox24*-specific, or a *vgf*-specific probe as indicated. Equal gel loading was confirmed by the hybridization with an 18 S-specific ribosomal RNA probe. These results are representative of three independent experiments.




FIG. 3. EGFR/ret stimulation of tyrosine phosphorylation of Snt in PC12. Cells were untreated (lanes 1 and 4) or treated with 100 ng/ml NGF (lanes 3 and 6) or EGF (lanes 2 and 5) for 5 min at 37 °C. Cell lysates were incubated with p13^{src}-agarose, eluted, and analyzed by immunoblot with anti-Tyr(P) (apTyr) antibody. The position of Snt is indicated.

ret^{Thr-918}, whereas stimulation of the mutant chimera caused its marked tyrosine phosphorylation (Fig. 5B) and overinduction of *krox24* gene expression (not shown).

DISCUSSION

Here we report data showing that the *ret* gene is able to differentiate the PC12 cells and that Ret carrying the MEN2B activating mutation is further inducible by ligand stimulation. To perform this study, we took advantage of an inducible system represented by a chimeric receptor (EGFR/ret) in which the tyrosine kinase activity of Ret was triggerable by EGF. When stimulated with EGF, PC12 cells transfected with EGFR/ret acquired a neuronal phenotype, characterized by long neuritic processes and the expression of immediate (*krox24*) as well as delayed (*vgf*) genes. Such phenotype is undistinguishable from that induced by NGF ("NGF phenotype"). This was further supported by the observation that the EGFR/ret chimera was also able to induce the tyrosine phosphorylation of Snt, a mol-

TABLE I
Transforming activity of the EGFR/ret^{Thr-918} chimera in NIH 3T3 cell fibroblasts

Transfected DNA	Transformation of NIH 3T3 cells ^a	
	–EGF	+EGF
FFU/pmol DNA		
EGFR/ret	1 × 10 ¹	1 × 10 ³
EGFR/ret ^{Thr-918}	1 × 10 ²	4 × 10 ³
LTR-3	<1 × 10 ¹	<1 × 10 ¹

^a Transfections were performed using 40 μ g of carrier calf thymus DNA. Where indicated, EGF (20 ng/ml) was added after 14 days. Focus-forming activity (in focus-forming units (FFU)) was scored at day 21 on duplicate plates transfected with 10-fold dilutions of the DNA of interest. Transforming activity is corrected for the efficiency of transfection calculated in parallel plates subjected to marker selection. Results are the means of three experiments performed in duplicate.




FIG. 4. Effects of stimulation of the EGFR/ret^{Thr-918} chimera on PC12 cell morphology. A phase contrast micrograph of ret^{Thr-918} (upper panels) or EGFR/ret^{Thr-918} (lower panels) PC12 cell transfectants is shown. A marker-selected mass population was grown for 72 h either in the absence or presence of EGF (100 ng/ml) as indicated. The same results were observed when four independent cell clones were analyzed for each cell line.

ecule that is regarded as a specific target of neurotrophic factors (23, 29).

The pattern of neuronal gene induction in PC12 cells, expressing the chronically active *ret*MEN2A and *ret*MEN2B alleles, is similar to that elicited by the acute stimulation of the EGFR/ret chimera. However, PC12-*ret*MEN2A and PC12-*ret*MEN2B cells displayed a less differentiated morphology with respect to EGF-stimulated PC12-EGFR/ret cells, since the former were characterized by a flat cell body and short neuritic processes ("MEN2 phenotype") (16) and the latter displayed a NGF phenotype. Whether or not these differences resulted from the kinetics of activation of the forms used, namely acute stimulation of EGFR/ret versus chronic activation of *ret*MEN2A and *ret*MEN2B, remains to be determined.

The inheritance of specific *ret* mutations causes distinct disease phenotypes, thus suggesting that some specific cell types undergo abnormal proliferation depending on the type of *ret* activation (via a MEN2A or via a MEN2B mutation) (2, 3). One possibility is that *ret*MEN2B activity could still be influenced by cell- or tissue-specific biological constraints, such as, for example, the density of the available ligand.

We thus investigated this possibility by using a mutated version of the chimera (EGFR/ret^{Thr-918}), harboring the MEN2B mutation. Consistent with the notion that MEN2B causes a gain of function of Ret, the EGFR/ret^{Thr-918} construct was able to transform NIH 3T3 cells and induce differentiation in PC12 cells, even in the absence of EGF. PC12 cells transfected with EGFR/ret^{Thr-918} showed a phenotype indistinguishable from PC12 transfected with *ret*MEN2B, thus confirming




FIG. 5. Stimulation of the EGFR/ret^{Thr-918} induces tyrosine phosphorylation of the receptor and mediates tyrosine phosphorylation of Snt in PC12 cells. **Panel A**, total cellular proteins from PC12 transfectants were immunoprecipitated (Ipt) with the anti-Ret polyclonal antibody (α ret) and analyzed by immunoblotting (blot) with the anti-Ret or anti-Tyr(P) (α pTyr) antibody, as indicated. Control untransfected cells (C), EGFR/ret (E/R), and EGFR/ret^{Thr-918} (E/R^{Thr-918}) transfectants were either untreated (–) or treated with 100 ng/ml EGF (+) for 5 min at 37 °C. Molecular mass markers are indicated in kilodaltons. The position of the EGFR/ret chimera is also indicated. **Panel B**, PC12 or PC12 cell transfectants were untreated (lanes 1, 4, and 6) or treated with 100 ng/ml EGF (lanes 2 and 5), or 100 ng/ml NGF (lane 3) for 5 min at 37 °C. Cell lysates were incubated with p13^{src}1-agarose, eluted, and analyzed by immunoblot with anti-Tyr(P) (α pTyr) antibody. The position of Snt is indicated.

that the expression in PC12 cells of a constitutive active Ret version results in a MEN2 phenotype. It is noteworthy that the MEN2B mutation was less effective in activating Ret function when cloned in the EGFR/ret construct with respect to the full-length ret. Since the difference between EGFR/ret and ret resides in their extracellular and transmembrane domains, it is likely that some specific characteristics of such domains confer to Ret this particular susceptibility to the activating effect of the MEN2B mutation.

However, despite the fact that the Met-918 to Thr mutation constitutively activates the chimera, the biological effects of the EGFR/ret^{Thr-918} construct were markedly sensitive to EGF triggering. EGF stimulation caused a marked increase of the transforming ability of the EGFR/ret^{Thr-918} construct and modified the phenotype of PC12-EGFR/ret^{Thr-918} cells, determining the overgrowth of long neuritic processes and a dramatic phosphorylation of Snt. These effects were consistent with the stim-

ulation of tyrosine phosphorylation of EGFR/ret^{Thr-918} caused by EGF.

These results show that the MEN2B mutation does not abrogate ligand responsiveness of Ret. However, we cannot discriminate whether the stimulation of the mutated Ret enhances the activity of the receptor, without changing the substrate specificity or, more likely, uncovers docking sites for new substrates. This ligand responsiveness may have important implications in the human diseases associated with retMEN2B mutations. It is likely that some of the differences in the disease phenotype between MEN2A and MEN2B syndromes could depend on the tissue distribution of the Ret ligand and on the different susceptibility of retMEN2A and retMEN2B alleles to the action of such a ligand.

REFERENCES

- Takahashi, M., Buma, Y., Iwamoto, T., Inaguma, Y., Ikeda, H., and Hiai, H. (1988) *Oncogene* 3, 571–578
- Mak, Y. F., and Ponder, B. A. J. (1996) *Curr. Opin. Genet. Dev.* 6, 82–86
- Goodfellow, P. J. (1994) *Curr. Opin. Genet. Dev.* 4, 446–452
- Mulligan, L. M., Kwok, J. B. J., Healey, C. S., Elsdon, M. J., Eng, C., Gardner, E., Love, D. R., Mole, S. E., Moore, J. K., Papi, L., Ponder, M. A., Telenius, H., Tunnacliffe, A., and Ponder, B. A. J. (1993) *Nature* 363, 458–460
- Donis-Keller, H., Dou, S., Chi, D., Carlson, K. M., Toshima, K., Lairmore, T. C., Howe, J. R., Moley, J. F., Goodfellow, P., and Wells, S. A., Jr. (1993) *Hum. Mol. Genet.* 2, 851–856
- Carlson, K. M., Dou, S., Chi, D., Scavarda, N., Toshima, K., Jackson, C. E., and Wells, S. A., Jr. (1994) *Proc. Natl. Acad. Sci. U. S. A.* 91, 1579–1583
- Hofstra, R. M. W., Landsvater, R. M., Ceccherini, I., Stulp, R. P., Stelwagen, T., Luo, Y., Pasini, B., Hoppener, J. W. M., van Amstel, H. K. P., Romeo, G., Lips, C. J. M., and Buys, C. H. C. M. (1994) *Nature* 367, 375–376
- Asai, N., Iwashita, T., Matsuyama, M., and Takahashi, M. (1995) *Mol. Cell. Biol.* 15, 1613–1619
- Santoro, M., Carlomagno, F., Romano, A., Bottaro, D. P., Dathan, N. A., Grieco, M., Fusco, A., Vecchio, G., Matoskova, B., Kraus, M. H., and Di Fiore, P. P. (1995) *Science* 267, 381–383
- Songyang, Z., Carraway, K. L., III, Eck, M. J., Harrison, S. C., Feldman, R. A., Mohammadi, M., Schlessinger, J., Hubbard, S. R., Smith, D. P., Eng, C., Lorenzo, M. J., Ponder, B. A. J., Mayer, B. J., and Cantley, L. C. (1995) *Nature* 373, 536–539
- Greene, L. A., and Tischler, A. S. (1976) *Proc. Natl. Acad. Sci. U. S. A.* 73, 2424–2428
- Marshall, C. J. (1995) *Cell* 80, 179–185
- Califano, D., Monaco, C., De Vita, G., D'Alessio, A., Dathan, N. A., Possenti, R., Vecchio, G., Fusco, A., Santoro, M., and de Francis, V. (1995) *Oncogene* 11, 107–112
- Pasini, B., Borrello, M. G., Greco, A., Bongarzone, I., Luo, Y., Mondellini, P., Alberti, L., Miranda, C., Arighi, E., Boccardi, R., Seri, M., Barone, V., Radice, M. T., Romeo, G., and Pierotti, M. A. (1995) *Nat. Genet.* 10, 35–40
- Borrello, M. G., Smith, D. P., Pasini, B., Bongarzone, I., Greco, A., Lorenzo, M. J., Arighi, E., Miranda, C., Eng, C., Alberti, L., Boccardi, R., Mondellini, P., Scopsi, L., Romeo, G., Ponder, B. A. J., and Pierotti, M. A. (1995) *Oncogene* 11, 2419–2427
- Califano, D., D'Alessio, A., Colucci-D'Amato, G. L., De Vita, G., Monaco, C., Santelli, G., Di Fiore, P. P., Vecchio, G., Fusco, A., Santoro, M., and de Francis, V. (1996) *Proc. Natl. Acad. Sci. U. S. A.* 93, 7933–7937
- Trupp, M., Arenas, E., Fainzilber, M., Nilsson, A. S., Sleber, B. A., Grigoriou, M., Kilkenny, C., Salazar-Gruoso, E., Pachnis, V., Arumae, U., Sariola, H., Saarma, M., and Ibanez, C. F. (1996) *Nature* 381, 785–789
- Durbec, P., Marcos-Gutierrez, C. V., Kilkenny, C., Grigoriou, M., Wartiovaara, K., Suvanto, P., Smith, D., Ponder, B., Costantini, F., Saarma, M., Sariola, H., and Pachnis, V. (1996) *Nature* 381, 789–793
- Jing, S., Wen, D., Yu, Y., Holst, P. L., Luo, Y., Fang, M., Tamir, R., Antonio, L., Hu, Z., Cupples, R., Louis, J. C., Hu, S., Altrock, B. W., and Fox, G. M. (1996) *Cell* 85, 1113–1124
- Treanor, J. J. S., Goodman, L., de Sauvage, F., Stone, D. M., Poulsen, K. T., Beck, C. D., Gray, C., Armanini, M. P., Pollock, R. A., Franz, H., Phillips, H. S., Goddard, A., Moore, M. W., Buj-Bello, A., Davies, A. M., Asai, N., Takahashi, M., Vanden, R., Henderson, C. E., and Rosenthal, A. (1996) *Nature* 382, 80–83
- Santoro, M., Wong, W. T., Aroca, P., Santos, E., Matoskova, B., Grieco, M., Fusco, A., and Di Fiore, P. P. (1994) *Mol. Cell. Biol.* 14, 663–675
- van Weering, D. H. J., Medema, J. P., van Puijnenbroek, A., Burgering, B. M. T., Baas, P. D., and Bos, J. L. (1996) *Oncogene* 11, 2207–2214
- Rabin, S. J., Cleghon, V., and Kaplan, D. R. (1993) *Mol. Cell. Biol.* 13, 2203–2213
- Chomczynski, P., and Sacchi, N. (1987) *Anal. Biochem.* 162, 156–159
- Possenti, R., Di Rocco, G., Nasi, S., and Levi, A. (1992) *Proc. Natl. Acad. Sci. U. S. A.* 89, 3815–3819
- Levi, A., and Alemà, S. (1991) *Annu. Rev. Pharmacol. Toxicol.* 31, 205–208
- Levi, A., Eldridge, J. D., and Paterson, B. M. (1985) *Science* 229, 393–395
- Milbrandt, J. (1987) *Science* 238, 797–799
- Peng, X., Greene, L. A., Kaplan, D. R., and Stephens, R. M. (1995) *Neuron* 15, 395–406

EXHIBIT D

Entrez Gene

Home
About
FAQ
Help
Gene Handbook
Statistics
Downloads
(FTP)

1: Ret

Ret oncogene [*Drosophila melanogaster*]

Other Aliases: Dmel_CG14396, CG1061, CG14396, D-ret, DRET, DmHD-59, MEN2, RET

Other Designations: Ret oncogene CG14396-PA, isoform A; Ret oncogene CG14396-PB, isoform B; Ret oncogene CG14396-PC, isoform C; Ret oncogene CG14396-PD, isoform D; Ret oncogene CG14396-PE, isoform E

Chromosome: 2L; **Location:** 39B1-39B1

GeneID: 43875

Mailing Lists

Gene
RefSeq

2: RET

Order cDNA clone, Links

Official Symbol RET and **Name:** ret proto-oncogene [*Homo sapiens*]

Other Aliases: CDHF12, HSCR1, MEN2A, MEN2B, MTC1, PTC, RET-ELE1, RET51

Other Designations: Hirschsprung disease 1; RET transforming sequence; cadherin family member 12; hydroxyaryl-protein kinase; multiple endocrine neoplasia and medullary thyroid carcinoma 1; oncogene RET; receptor tyrosine kinase; ret proto-oncogene (multiple endocrine neoplasia and medullary thyroid carcinoma 1, Hirschsprung disease)

Chromosome: 10; **Location:** 10q11.2

Annotation: Chromosome 10, NC_000010.9 (42892533..42944955)

MIM: 164761

GeneID: 5979

3: Ret

Links

Official Symbol Ret and **Name:** ret proto-oncogene [*Rattus norvegicus*]

Other Designations: Ret gene for receptor tyrosin; Ret proto-oncogene (multiple endocrine neoplasia MEN2A MEN2B and medullary thyroid carcinoma 1 Hirschsprung disease); receptor tyrosine kinase

Chromosome: 4; **Location:** 4q42

Annotation: Chromosome 4, NC_005103.2 (154448179..1544491103, complement)

GeneID: 24716

4: RET

Links

ret proto-oncogene [*Canis lupus familiaris*]

Other Designations: ret proto-oncogene (multiple endocrine neoplasia and medullary thyroid carcinoma 1, Hirschsprung disease); ret tyrosine kinase

Chromosome: 28

Resources

Related Sites

BLAST
Entrez Genome
Genome
Projects
Genomic
Biology
GEO
HomoloGene
Map Viewer
OMIM
Probe
RefSeq
UniGene
UniSTS

NIH cDNA
Tutorial

Entrez

Global Search
About | Help

NCBI

Home
Site Map
Search
Handbook

5: Ret

Official Symbol Ret and Name: ret proto-oncogene [*Mus musculus*]

Other Aliases: PTC, RET51, RET9, c-Ret

Chromosome: 6; **Location:** 6 53.2 cM

Annotation: Chromosome 6, NC_000072.5 (118101766..118147762, complement)

GeneID: 19713

Order cDNA clone, Links

6: RET

ret proto-oncogene [*Gallus gallus*]

Chromosome: 6

Annotation: Chromosome 6, NC_006093.2 (5877654..5953679, complement)

GeneID: 396107

Links

7: RET

ret proto-oncogene [*Pan troglodytes*]

GeneID: 449594

Links

8: LOC450415

similar to ret proto-oncogene isoform c; hydroxyaryl-protein kinase; RET transforming sequence; oncogene RET; cadherin family member 12 [*Pan troglodytes*]

Chromosome: 10

GeneID: 450415

This record was discontinued.

Links

9: ret

reticulated [*Drosophila melanogaster*]

Other Aliases: FBgn0010168

Chromosome: 1; **Location:** 1-

GeneID: 44319

This record was discontinued.

10: RFPL4A

Official Symbol RFPL4A and Name: ret finger protein-like 4A [*Homo sapiens*]

Links

EXHIBIT E

1: RET ret proto-oncogene [Homo sapiens]

GeneID: 5979

updated 16-Sep-2007


Summary

Official Symbol	RET	provided by HGNC
Official Full Name	ret proto-oncogene	provided by HGNC
Primary source	HGNC:9967	
See related	Ensembl:ENSG00000165731; HPRD:01266; MIM:164761	
Gene type	protein coding	
RefSeq status	Reviewed	
Organism	Homo sapiens	
Lineage	Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini; Catarrhini; Hominidae; Homo	
Also known as	PTC; MTC1; HSCR1; MEN2A; MEN2B; RET51; CDHF12; RET-ELE1	

Summary This gene, a member of the cadherin superfamily, encodes one of the receptor tyrosine kinases, which are cell-surface molecules that transduce signals for cell growth and differentiation. This gene plays a crucial role in neural crest development, and it can undergo oncogenic activation in vivo and in vitro by cytogenetic rearrangement. Mutations in this gene are associated with the disorders multiple endocrine neoplasia, type IIA, multiple endocrine neoplasia, type IIB, Hirschsprung disease, and medullary thyroid carcinoma. Two transcript variants encoding different isoforms have been found for this gene. Additional transcript variants have been described but their biological validity has not been confirmed.

Genomic regions, transcripts, and products

Go to reference sequence details



Genomic context

chromosome: 10; Location: 10q11.2

See RET in MapViewer



Bibliography

Related Articles in PubMed

PubMed links

GeneRIFs: Gene References Into Function

- 1.These data highlight the pivotal role of the RET gene in both isolated and syndromic Hirschsprung disease.
- 2.Of these 86 variations, 8 proved to be in regions highly conserved among different vertebrates and within putative transcription factor binding sites. We therefore considered these as candidate disease-associated variants.
- 3.frequency of RET rearrangements in papillary thyroid carcinoma for the Polish population
- 4.our results suggest that SPRY2 regulates GDNF-dependent proliferation and differentiation of TGW neuroblastoma cells mediated by RET tyrosine kinase.
- 5.RET/PTC rearrangement in thyroid tumors. Review
- 6.Point mutation in exon 14 at codon 804 of the RET proto-oncogene locus in a case of lymph node metastases of medullary and papillary thyroid carcinoma.
- 7.High prevalence of BRAF mutations in thyroid cancer is genetic evidence for constitutive activation of th RET/PTC-RAS-BRAF signaling pathway in papillary thyroid carcinoma.
- 8.The Cys630 RET genotype may have a more vigorous transforming activity than currently thought and

can cause medullary thyroid carcinoma in RET gene carriers within the first year of life.

9. RET, a receptor tyrosine kinase involved with differentiation, was consistently up-regulated throughout the time course of retinoic acid treatment in the majority of neuroblastic tumor cell lines.
10. Findings identify RET as a novel substrate of PTPRJ and suggest that PTPRJ expression levels may affect tumor phenotype associated with RET/PTC1 and RET(C634R) mutants.
11. Copy gain of PDGFB occurs in a subset of tumors showing no evidence of mutated BRAF or rearranged ret, suggesting that copy gain of PDGFB may underlie the increased expression of platelet-derived growth factor described recently in the literature.
12. Mutations of RET proto-oncogene may play an important role in the pathogenesis of Chinese patients with Hirschsprung disease.
13. RET gene mutation may explain the wide clinical variability associated with germline mutations at codon 804 in medullary thyroid carcinoma/multiple endocrine neoplasia type 2A patients.
14. The RET/PTC1 oncogene activates a proinflammatory program, provide a direct link between a transforming human oncogene, inflammation, and malignant behavior.
15. Mutagenesis analysis revealed that Tyr981 within the intracellular domain of RET was crucial for the interaction with SH2-Bbeta. Morphological evidence showed that SH2-Bbeta and RET colocalized in mesencephalic neurons.
16. Germ-Line Mutation in RET proto-oncogene is associated with Multiple Endocrine Neoplasia
17. Three new somatic cell missense mutations of the RET proto-oncogene associated with sporadic medullary thyroid carcinoma (MTC).
18. C630R mirrors C634R in penetrance and in early age of onset of medullary thyroid carcinoma
19. RET rearrangements may not play any distinctive role in driving histotype development and cancer progression in papillary thyroid carcinomas.
20. In RET mutation carriers in Hirschsprung's disease, the gut caliber change was almost identical to the histologic transition in cases of short segment aganglionosis, whereas these were markedly dissociated in cases exhibiting extensive aganglionosis
21. In the presence of RET oncoproteins, both RAI and GAB 1 are tyrosine-phosphorylated, and the stoichiometry of this interaction remarkably increases

22. Analysis of the RET gene revealed neither linkage nor mutations in Hirschsprung's disease mapping.
23. new missense point mutation in exon 8 of the RET gene (1597G-->T) corresponding to a Gly(533)Cys substitution in the cysteine-rich domain of RET protein
24. Not only RET mutations but also RET polymorphic variants may contribute to the occurrence of total intestinal aganglionosis.
25. Single nucleotide polymorphisms in the RET oncogene may play a role in sporadic papillary thyroid carcinoma.
26. A founding locus within the RET proto-oncogene may account for a large proportion of apparently sporadic Hirschsprung disease and a subset of cases of sporadic medullary thyroid carcinoma
27. Identification of a heterozygous germ line missense mutation at codon 634 of exon 11 in the RET gene that causes a cysteine to arginine amino acid substitution in a MEN2A patient.
28. Dissecting Hirschsprung disease. RET is the main gene conferring susceptibility.
29. Segregation at three loci explains familial and population risk in Hirschsprung disease. We show oligogenic inheritance of S-HSCR, the 3p21 and 19q12 loci as RET-dependent modifiers, and a parent-of-origin effect at RET.
30. Patients with RET codon 790/791 mutations seemed to have a less aggressive clinical course compared with patients with classic multiple endocrine neoplasia type 2A syndrome.
31. germline mutation of the RET proto-oncogene in members of Slovak families with multiple endocrine neoplasia 2
32. Single nucleotide polymorphism in ret is associated with the aggressive growth of pancreatic cancers
33. Amplification and overexpression of mutant RET in multiple endocrine neoplasia type 2-associated medullary thyroid carcinoma. RET germline mutation in codon 634. Tandem duplication. Genomic chromosome 10 abnormalities increase mutant RET mRNA.
34. Expression of the mitogenic and invasive phenotype of RET/PTC-transformed thyroid cells is stimulated by osteopontin.
35. Data report the crystal structure of GFRalpha1 domain 3, and the effects of specific mutations on GDNF binding and RET phosphorylation.

36. RET/PTC expression phosphorylates the Y701 residue of STAT1, a type II interferon (IFN)-responsive protein.
37. the absence of RET alterations in all cases of C-cell hyperplasia
38. Loss-of-function germline mutations of the RET proto-oncogene are reported in familial and sporadic cases of Hirschsprung disease (HSCR) with a variable frequency
39. Selective disruption of oncogenic RET signaling in medullary thyroid carcinoma in vitro and in vivo is associated with loss of the neoplastic phenotype of medullary thyroid carcinoma.
40. RET expression leads to increased HSF1 activation, which correlates with increased expression of stress response genes. RET may be directly responsible for expression of stress response proteins and the initiation of stress response.
41. RET has roles in neoplastic transformation [review]
42. Papillary carcinomas with high RET/PTC1 expression showed an association trend for large tumor size.
43. direct interaction between RET and a broad range of effector molecules that may contribute to tumor pathogenesis
44. gene expression impairment seems to be at the basis of the association of HSCR disease with several RET polymorphisms, allowing us to define a predisposing haplotype spanning from the promoter to exon 2.
45. mutations of the RET protooncogene were analyzed in Russian patients with inherited or sporadic medullary thyroid carcinoma. The most common mutation affected codon 918 to cause substitution of methionine with threonine and accounted for 31.6% alleles.
46. The finding of a somatic deletion in RET exon 15 clarified the sporadic nature of a medullary thyroid carcinoma suspected to be familial. A 12 bp deletion within the catalytic domain of the protooncogene RET.
47. Possible pathogenesis of papillary thyroid carcinoma caused by exon 13 and 14 RET mutations that affect the intracellular domain of ret proto-oncogene protein.
48. Hirschsprung associated GDNF mutations do not prevent RET activation
49. Koreans showed increased RET gene expression in papillary thyroid carcinoma.

50. Nuclease-resistant aptamers that recognize the human receptor tyrosine kinase RET were obtained using RET-expressing cells as targets in a modified random pool of sequences.
51. RET point mutants for follicular thyroid cells may account for the occurrence of papillary thyroid carcinoma in patients affected by familial medullary thyroid carcinoma
52. RET signals through focal adhesion kinase in medullary thyroid cancer cells.
53. RET/PTC is able to phosphorylate the Y315 residue of PKB, an event that results in maximal activation of PKB for RET/PTC-induced thyroid tumorigenesis.
54. Possibility that lower-penetrance RET mutations may contribute to the list of causes of familial pheochromocytomas
55. Germline mutation on the RET gene was present in patients with pheochromocytoma or functional paraganglioma.
56. Two Hirschsprung disease-associated haplotypes derive from a single founding locus, extending up to intron 19 and successively rearranged in correspondence with a high recombination rate region located between the proximal and distal end of the RET gene.
57. These results reveal novel roles of key RET-dependent signaling pathways in embryonic kidney development and provide murine models and new insights into the molecular basis for CAKUT.
58. Mutational screening of RET revealed 9 different mutations, present in 26 of the 114 MEN 2 Spanish patients.
59. Timing and extent of prophylactic thyroidectomy can be modified by individual RET mutation
60. RET dysfunction has a crucial role and discusses RET as a potential therapeutic target.
61. These results suggest that RFP is a mediator connecting several MBD proteins and allowing the formation of a more potent transcriptional repressor complex.
62. Children of families with RET cysteine mutations may develop early metastatic medullary carcinoma of the thyroid gland.
63. Study demonstrates that the interaction between RET and PHOX2B polymorphisms has a substantial impact on risk of Hirschsprung's disease
64. The BRAF(V599E) mutation appears to be an alternative event to RET/PTC rearrangement rather than t

RAS mutations, which are rare in PTC. BRAF(V599E) may represent an alternative pathway to oncogenic MAPK activation in PTCs without RET/PTC activation.

65. Letter discussing RET mutations in distinguishing between sporadic and familial medullary thyroid carcinoma.
66. the RET proto-oncogene mutation Y791F, characterized by a low penetrance, occurs comparatively frequently among patients with normal serum calcitonin concentrations
67. if the association between Hashimoto's thyroiditis and thyroid cancer exists, its molecular basis is different from RET/PTC rearrangement
68. Specific nucleotide and amino acid exchanges at codon 634 might have a direct impact on tumor aggressiveness in MEN 2A.
69. All Ret dominant-negative/+ mice died by 1 month of age and had distal intestinal aganglionosis reminiscent of Hirschsprung disease (HSCR) in humans
70. Mutations in medullary thyroid cancer and in multiple endocrine neoplasia 2.
71. The newly identified RET/N777S germline mutation is a low-penetrant cause of medullary thyroid carcinoma.
72. Medullary thyroid carcinoma as part of multiple endocrine neoplasia type 2 in a family with a mutation in RET proto oncogene.
73. results indicate a possible association between the presence of lymph node involvement at the time of diagnosis (extent of disease) of medullary thyroid carcinoma and L769L or S836S polymorphism.
74. Differentiation of cardiac ganglionic cells is affected, after RETINOIC ACID treatment, by the down-regulation of c-Ret.
75. RET requires coupling of Gab1 to phosphatidylinositol 3-kinase for function in human tumor cells
76. Expression of a human Ret proto-oncogene with the MEN 2B mutation does not cause any features of MEN 2B in mice.
77. RET/PTC and CK19 have roles in progression of papillary thyroid carcinoma
78. Data show that the RET receptor (RET/PTC), Ras and BRAF function along a linear oncogenic signaling cascade in which RET/PTC induces RAS-dependent BRAF activation and RAS- and BRAF-

dependent ERK activation.

- 79.early detection of RET proto-oncogene mutation is crucial for prevention of thyroidectomy in multiple endocrine neoplasia type 2 children
- 80.Y1062 is a critical regulator of Ret9 signaling and suggest that Ret51-specific motifs are likely to inhibit the activity of this isoform
- 81.new missense point mutation in exon 5 in familial medullary thyroid carcinoma
- 82.Ret tyrosine 981 constitutes the major binding site of the Src homology 2 domain of Src and therefore the primary residue responsible for Src activation upon Ret engagement
- 83.With this study we excluded influence of the G691S polymorphism on RET mRNA expression, development of somatic RET mutation, the linkage with germline RET mutation, younger onset of medullary thyroid carcinoma, and clinical outcome of the disease.
- 84.RET activation closely parallels the morphological changes, that it is restricted to those areas of the tumor with the cytological alterations and that it is detectable in both mono- and polyclonal tumors
- 85.Shp2 activity required for RetM918T-induced Akt activation. Shp2 downstream mediator of mutated receptors RetC634Y and RetM918T. Shp2 acts as limiting factor in Ret-associated endocrine tumors, in neoplastic syndromes multiple endocrine neoplasia.
- 86.Persephin/GFRalpha4 is unable to recruit RET protein into lipid rafts.
- 87.Single nucleotide polymorphisms of RET is associated with Hirschsprung disease
- 88.A novel Val648Ile substitution in RET protooncogene observed in a Cys634Arg multiple endocrine neoplasia type 2A kindred presenting with an adrenocorticotropin-producing pheochromocytoma.
- 89.Substantial discrimination between predicted functional classes of RET mutations and disease severity even for a multigenic disease such as Hirschsprung disease.
- 90.The histone acetylation level was evaluated by the chromatin immunoprecipitation method applied to cells displaying different degrees of endogenous RET expression.
- 91.evaluation of noncoding sequences at the zebrafish ret locus conserved among teleosts, and at the human RET locus, conserved among mammals
- 92.Cell type-specific functions involve a competitive recruitment of different phosphotyrosine binding

adaptor molecules by RET that activate selective signaling pathways.

93. Ret mutations in thyroid tumorigenesis.

94. tumor samples from FMTC patients showed strong nuclear staining of phosphorylated ERK1/2 and Ser (727) STAT3; FMTC-RET mutants activate a Ras/ERK1/2/STAT3 Ser(727) pathway, which plays an important role in cell mitogenicity and transformation.

95. A germline RET mutation at codon 603 in exon 10 is associated with both medullary and nonmedullary thyroid cancer in a kindred.

96. there is a low-penetrance pheochromocytoma susceptibility locus in a region upstream of the putative loci for Hirschprung disease and apparently sporadic thyroid carcinoma.

97. RET oligonucleotide microarray for the detection of RET mutations in multiple endocrine neoplasia type syndromes

98. 5'-End RET splicing: absence of variants in normal tissues and intron retention in pheochromocytomas.

99. Dok-6 binds to the phosphorylated Ret Tyr(1062) residue resulting in phosphorylation of tyrosine residue(s) located within the unique C terminus of Dok-6 predominantly through a Src-dependent mechanism

100. relationship between RET oncogene and Chinese patients with Hirschsprung's disease

101. mutated in papillary thyroid cancer.

102. These observations lend support to the idea that both RET alleles have a role in pathogenesis of Hirschsprung's disease, in a dose-dependent fashion. We also showed that the c135G/A polymorphism modifies the phenotype.

103. determination of mutation specific gene expression profiles in papillary thyroid carcinoma

104. Erk8 has a role as a novel effector of RET/PTC3 and, therefore, RET biological functions

105. First molecular studies on a complex germline RET mutation lying in the juxtamembrane region of the receptor are reported in medullary thyroid carcinoma

106. The molecular basis for HPT has been further elucidated by the detection of inactivating germline mutations in the CaSR gene in familial hypocalciuric hypercalcemia syndrome and in the RET genes in the familial forms of HPT.

107. These findings provide evidence for a novel cooperative interaction between VEGFR2 and RET that mediates VEGF-A functions in ureteric bud cells.
108. RET codon 691 polymorphism is associated with radiation induced tumors with a C-cell hyperplasia of thyroid tumors
109. role in regulating rac activity and lamellipodia formation
110. genetic interaction between mutations in RET and EDNRB is an underlying mechanism for Hirschsprung disease
111. RET expression in papillary thyroid cancer from patients irradiated in childhood for benign conditions.
112. family in which the MEN 2A and the HSCR phenotypes are associated with a single point mutation in exon 10 of the RET proto-oncogene. polymorphic sequence variants of the RET proto-oncogene.
113. a protective role of this low-penetrant haplotype in the pathogenesis of HSCR and demonstrate a possible functional effect linked to RET messenger RNA expression.
114. analysis of RET somatic mutations supports the differentiation between sporadic and inherited medullary thyroid carcinoma
115. Analysis of mutation of protooncogene RET are presented in patients with thyroid medullary carcinoma
116. Four novel intronic mutations that have a strong association with the HSCR phenotype were identified in Hirschsprung disease patients
117. the G691S and S904S variants of RET may somehow play a role on the age of onset of MEN 2A
118. PHOX2A, but not PHOX2B, seems to act directly on the c-RET promoter
119. RET/PTC associates with STAT3 and activates it by the specific phosphorylation of the tyrosine 705 residue. STAT3 activation by the RET/PTC tyrosine kinase is one of the critical signaling pathways for the regulation of specific genes.
120. medullary thyroid carcinoma manifested in new RET mutation and RET polymorphism.
121. Familial medullary thyroid carcinoma: clinical variability and low aggressiveness associated with RET mutation at codon 804.
122. Activation of RET tyrosine kinase regulates interleukin-8 production by multiple signaling pathways

123. A deletion of the chromosomal region including the RET proto-oncogene is involved in the pathogenesis of SCLC
124. Mass spectrometric analysis revealed that RET Tyr(806), Tyr(809), Tyr(900), Tyr(905), Tyr(981), Tyr(1062), Tyr(1090), and Tyr(1096) were autophosphorylation sites.
125. RET enhancer modulates expression in the enteric nervous system consistent with its proposed role in Hirschsprung disease
126. Significant association of the S691 allele with medullary thyroid carcinoma
127. Characterization of the R833C substitution suggests that this tyrosine kinase mutation confers a weak activating potential upon RET but introduces an intracellular cysteine which activates RET.
128. Strong propensity to self-association in the RET-transmembrane underlies - and may be required for - dimer formation and oncogenic activation of juxtamembrane cysteine mutants of RET
129. Association of high-level Ret proto-oncogene protein expression with neuronal morphology suggests that the variable overexpression of Ret in pheochromocytomas might in part be an epiphenomenon, reflecting the known phenotypic plasticity of these tumors
130. Ret expression is significantly higher in thyroid papillary carcinoma than benign thyroid tissue; and this characteristic can have important diagnostic value.
131. Association of RET IVS1-126G>T variant with sporadic medullary thyroid cancer in a cohort of 104 patients
132. Mechanisms leading to RET oncogenic conversion
133. Findings support the notion that both RET alleles are involved in the pathogenesis of a subgroup of Hirschsprung disease patients in a dose-dependent fashion
134. These findings establish a mechanism for the differential down-regulation of RET9 and RET51 signaling that could explain the apparently paradoxical activities of these two RET isoforms.
135. A RET haplotype (A-C-A) composed of alleles at three SNPs is associated with reduced RET gene expression in Hirschsprung patients.
136. A common non-coding RET variant within a conserved enhancer-like sequence in intron 1 is significantly associated with Hirschsprung disease susceptibility

- 137. Gas1 is related to the GDNF alpha receptors and regulates Ret signaling
- 138. We show that RA-induced differentiation is mediated by a positive autocrine loop that sustains Ret downstream signaling and depends on glial cell-derived neurotrophic factor expression and release.
- 139. the RET finger protein has a role in estrogen receptor-mediated transcription in tumor cells
- 140. A variant located in the 3' untranslated region of the RET gene, which slows down mRNA decay in patients with Hirschsprung disease.
- 141. study demonstrated RET amplification in all 3 cases of radiation-associated thyroid cancers (papillary thyroid cancer (PTC) & anaplastic thyroid cancer (ATC) but not in sporadic well-differentiated PTC; RET amplification was observed in all 6 cases of ATCs
- 142. critical role of the immunoglobulin domain in regulation of the localization of human PTPmu in bovine cells
- 143. The presence of coiled-coil domains in the ktn1/ret fusion protein (PTC8) suggests ligand-independent dimerization and thus constitutive activation of the ret tyrosine kinase domain.

Interactions

Description				
Product	Interactant	Other Gene	Complex	Source
RET interacts with FRS2				
NP_065681.1	NP_006645.2	FRS2	BIND	PubMed
RET9 phosphorylates PLC-gamma via its amino-terminal SH2 domain. This interaction was modelled on a demonstrated interaction between RET9 from human and PLC-gamma from cow.				
NP_065681.1	NP_002651.2	PLCG1	BIND	PubMed
RET9 phosphorylates Shc.				
NP_065681.1	NP_003020.2	SHC1	BIND	PubMed
NP_066124.1	NP_005179.1	CBL	HPRD	PubMed
NP_066124.1	Docking protein 1	DOK1	HPRD	PubMed
NP_066124.1	DOK2	DOK2	HPRD	PubMed
NP_066124.1	Docking protein 3	DOK3	HPRD	PubMed

General gene information

Markers

STS-T15350(e-PCR)

Links: UniSTS:15110

Alternate names: RH40273; sts-T15350

RH66458(e-PCR)

Links: UniSTS:20244

Alternate name: stSG35889

D10S1566(e-PCR)

Links: UniSTS:23844

Alternate names: G00-588-355; GDB:588355; RH50947; UTR-03097; WI-7098

G59921(e-PCR)
Links: UniSTS:137202
Alternate name: SHGC-130704

GDB:193843(e-PCR)
Links: UniSTS:155759

GDB:281473(e-PCR)
Links: UniSTS:156383

GDB:342173(e-PCR)
Links: UniSTS:156640

GDB:342177(e-PCR)
Links: UniSTS:156641

GDB:574049(e-PCR)
Links: UniSTS:157789

GDB:579598(e-PCR)
Links: UniSTS:157821

GDB:580702(e-PCR)
Links: UniSTS:157848

GDB:592843(e-PCR)
Links: UniSTS:157925

PMC23476P2(e-PCR)
Links: UniSTS:272204

Genotypes

See RET SNP GeneView Report
See RET SNP Genotype Report

Phenotypes

Central hypoventilation syndrome, congenital
MIM: 209880
Colonic aganglionosis, total, with small bowel involvement
MIM: 164761
Hirschsprung disease

- MIM: 142623
Medullary thyroid carcinoma
- MIM: 155240
Multiple endocrine neoplasia IIA
- MIM: 171400
Multiple endocrine neoplasia IIB
- MIM: 162300
Pheochromocytoma
- MIM: 171300

Pathways

KEGG pathway: Thyroid cancer
05216

Homology

Mouse, Rat
Map Viewer

GeneOntology

Provided by GOA

Function	Evidence
ATP binding	IEA
ATP binding	NAS
calcium ion binding	IEA
kinase activity	IEA
nucleotide binding	IEA
protein-tyrosine kinase activity	NAS
protein-tyrosine kinase activity	TAS
PubMed 7824936	
receptor activity	IEA
receptor activity	TAS
PubMed 7824936	

transferase activity	IEA
----------------------	-----

Process	Evidence
homophilic cell adhesion	IEA
posterior midgut development	TAS
PubMed 8114939	
protein amino acid phosphorylation	NAS
protein amino acid phosphorylation	TAS
PubMed 7824936	
signal transduction	TAS
PubMed 7824936	

Component	Evidence
integral to membrane	IEA
membrane	IEA

General protein information

Names

ret proto-oncogene
oncogene RET
Hirschsprung disease 1
receptor tyrosine kinase
RET transforming sequence
cadherin family member 12
hydroxyaryl-protein kinase
multiple endocrine neoplasia and medullary thyroid carcinoma 1
ret proto-oncogene (multiple endocrine neoplasia and medullary thyroid carcinoma 1, Hirschsprung disease)

NP_065681.1

EC 2.7.10.1

NP_066124.1
EC 2.7.10.1

NCBI Reference Sequences (RefSeq)

RefSeqs maintained independently of Annotated Genomes

These reference sequences exist independently of genome builds.

mRNA and Protein(s)

1. NM_020630.4-NP_065681.1 ret proto-oncogene isoform c

Description	Transcript Variant: This variant (4) differs in the 3' UTR and coding region compared to variant 2. The resulting isoform (c) is shorter and has a distinct C-terminus compared to isoform a. This isoform is also known as Ret9.
Source sequence(s) Conserved Domains (10) summary	AI472270,BC003072,BC004257,BE261914,BM703293,DA100452,DA911581,X12949
	cd00031 Location:173-358 Blast Score:188 Location:178-302 Blast Score:188 Location:43-264 Blast Score:106 Location:43-245 Blast Score:106
	CA; Cadherin repeat domain; Cadherins are glycoproteins involved in Ca ²⁺ -mediated cell adhesion; these domains occur as repeats in the extracellular regions which are thought to mediate cell-cell contact when bound to calcium; plays a role in cell fate,
	cd00192 Location:716-1008 Blast Score:971 Location:872-881 Blast Score:971 Location:788-893 Blast Score:971 Location:892-921 Blast Score:971 Location:813-915 Blast Score:971
	TyrKc; Tyrosine kinase, catalytic domain. Phosphotransferases; tyrosine-specific kinase subfamily.
	smart00220 Location:726-1004 Blast Score:449
	S_TKc; Serine/Threonine protein kinases, catalytic domain; Phosphotransferases. Serine or threonine-specific kinase subfamily.

2. NM_020975.4-NP_066124.1 ret proto-oncogene isoform a

Description	Transcript Variant: This variant (2) represents the longer transcript and encodes the longer isoform (a). This isoform is also known as Ret51.
Source sequence(s) Consensus CDS	AC010864,BC003072,BC004257,BM661773,DA100452,X12949 CCDS7200.1

Conserved Domains (10) summary	
cd00031 Location:173-358 Blast Score:188 Location:178-302 Blast Score:188 Location:43-264 Blast Score:105 Location:43-245 Blast Score:105	CA; Cadherin repeat domain; Cadherins are glycoproteins involved in Ca2+-mediated cell-cell adhesion; these domains occur as repeats in the extracellular regions which are thought to mediate cell-cell contact when bound to calcium; plays a role in cell fate,
cd00192 Location:716-1008 Blast Score:973 Location:872-881 Blast Score:973 Location:788-893 Blast Score:973 Location:892-921 Blast Score:973 Location:813-915 Blast Score:973	TyrKc; Tyrosine kinase, catalytic domain. Phosphotransferases; tyrosine-specific kinase subfamily.
smart00220 Location:726-1004 Blast Score:451	S_TKc; Serine/Threonine protein kinases, catalytic domain; Phosphotransferases. Serine or threonine-specific kinase subfamily.

RefSeqs of Annotated Genomes: Build 36.2

The following sections contain reference sequences that belong to a specific genome build.

Reference assembly

Genomic

1. NC_000010.9 Reference assembly

Range	Download
42892533..42944955	GenBank FASTA

2. NT_033985.6

Range	Download
975840..1028262	GenBank FASTA

Alternate assembly (based on Celera assembly)

Genomic

1.	AC_000053.1 Alternate assembly (based on Celera assembly)	
	Range	39575673..39628104
	Download	GenBank FASTA
2.	NW_924606.1	
	Range	986821..1039252
	Download	GenBank FASTA

Suppressed Reference Sequence(s)

The following Reference Sequences have been suppressed. Explain

1.	NM_000323.2: Suppressed sequence
Description	NM_000323.2 was permanently suppressed because currently not enough support exists for the transcript and the protein.
2.	NM_020629.2: Suppressed sequence
Description	NM_020629.2 was permanently suppressed because currently not enough support exists for the transcript and the protein.

Related Sequences

Nucleotide	Protein
Genomic AC010864.11 (75336..127758)	None
Genomic AF032124.1	AAB97168.1
Genomic AF520975.1	AAM77275.1
Genomic AF520979.1	AAM77279.1
Genomic AF520983.1	AAM77283.1
Genomic AJ243297.1	CAB46483.1
Genomic CH471160.1	EAW86576.1
	EAW86577.1
	EAW86578.1
	EAW86579.1
	EAW86580.1

Genomic	D00617.1	EAW86581.1
Genomic	S80097.1	EAW86582.1
		EAW86583.1
		EAW86584.1
		None
		AAB47046.1
		AAB47047.1
		AAB47046.1
		AAB47047.1
		AAB50647.2
		AAD14423.1
		CAA75753.1
		None
		None
		AAH03072.1
		AAH04257.1
		None
		None
		None
		None
		None
		None
		None
		None
		None
		None
		CAA33333.1
		CAA73131.1

Protein Accession	Links
O43519	GenPept UniProtKB/TrEMBL
P07949	GenPept UniProtKB/Swiss-Prot
Q15850	GenPept UniProtKB/TrEMBL
Q2VJ45	GenPept UniProtKB/TrEMBL

Q8IZR8	GenPept	UniProtKB/TrEMBL
Q8NFE8	GenPept	UniProtKB/TrEMBL
Q99886	GenPept	UniProtKB/TrEMBL
Q9BTB0	GenPept	UniProtKB/TrEMBL
Q9BTX6	GenPept	UniProtKB/TrEMBL
Q9UE13	GenPept	UniProtKB/TrEMBL
Q9UM84	GenPept	UniProtKB/TrEMBL
Q9UM90	GenPept	UniProtKB/TrEMBL
Q9UMQ4	GenPept	UniProtKB/TrEMBL
Q9UQV8	GenPept	UniProtKB/TrEMBL

Additional Links

- MIM 164761
- GeneTests for MIM: 164761
- HPRD 01266
- UniGene Hs.350321

Display

Full Report

Show

20

Send to


Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Privacy Statement | Freedom of Information Act | Disclaimer

EXHIBIT F

CANCER

*Principles & Practice
of Oncology*

6th Edition



www.LWWoncology.com



LIPPINCOTT WILLIAMS & WILKINS

A Wolters Kluwer Company

Philadelphia • Baltimore • New York • London
Buenos Aires • Hong Kong • Sydney • Tokyo

Acquisitions Editor: J. Stuart Freeman, Jr.

Development Editors: Susan Rhyner, Managing Editor

Anne Snyder, Senior Development Editor

Stephanie Harris, Associate Editor

Supervising Editor: Toni Ann Scaramuzzo

Production Editors: Kim Langford and Shannon Garza, Silverchair Science + Communications

Manufacturing Manager: Tim Reynolds

Compositor: Silverchair Science + Communications

Printer: Quebecor World, Taunton, MA

6th edition

© 2001 by LIPPINCOTT WILLIAMS & WILKINS

530 Walnut Street

Philadelphia, PA 19106 USA

LWW.com

Copyright © 1993, 1989, 1985, 1982 by J.B. Lippincott Company. All rights reserved. This book is protected by copyright. No part of this book may be reproduced in any form or by any means, including photocopying, or utilized by any information storage and retrieval system without written permission from the copyright owner, except for brief quotations embodied in critical articles and reviews. Materials appearing in this book prepared by individuals as part of their official duties as U.S. government employees are not covered by the above-mentioned copyright.

Printed in the USA

Library of Congress Cataloging-in-Publication Data

Library of Congress Control Number: 89-649-721

Cancer: principles and practice of oncology [edited by] Vincent T. DeVita, Jr., Samuel Hellman, Steven A. Rosenberg; 319 contributors.—6th

ISSN 0892-0567

ISBN 0-781-72387-6

Care has been taken to confirm the accuracy of the information presented and to describe generally accepted practices. However, the authors, editors, and publisher are not responsible for errors or omissions or for any consequences from application of the information in this book and make no warranty, expressed or implied, with respect to the currency, completeness, or accuracy of the contents of the publication. Application of this information in a particular situation remains the professional responsibility of the practitioner.

The authors, editors, and publisher have exerted every effort to ensure that drug selection and dosage set forth in this text are in accordance with current recommendations and practice at the time of publication. However, in view of ongoing research, changes in government regulations, and the constant flow of information relating to drug therapy and drug reactions, the reader is urged to check the package insert for each drug for any change in indications and dosage and for added warnings and precautions. This is particularly important when the recommended agent is a new or infrequently employed drug.

Some drugs and medical devices presented in this publication have Food and Drug Administration (FDA) clearance for limited use in restricted research settings. It is the responsibility of health care providers to ascertain the FDA status of each drug or device planned for use in their clinical practice.

10 9 8 7 6 5 4 3 2 1

- Respiratory disorders (*continued*)
 mechanism of lung injury in, 2897-2898
 from methotrexate, 392
 in opioid therapy, 2997
 pulmonary function tests in, 2901
 risk factors in, 2898, 2898t
 signs and symptoms in, 2898-2901
 in testicular germ cell tumors, 1512
 treatment of, 2901
 hospice care in, 3081-3082, 3084
 infections
 in aspergillosis, 2827, 2827f, 2828, 2853-2854
 computed tomography in, 667, 2827f, 2828
 Pneumocystis carinii, 2839
 pulmonary infiltrates in, 2851-2854, 2852t
 differential diagnosis in, 2851, 2852t
 viral, 2837-2838
 laparoscopy in, 740
 in lung cancer, 932
 palliative care in, 974
 postoperative, 944
 preoperative assessment of, 944
 and radiation therapy, 946, 995
 prevention of, 2894
 small cell, 985
 in pleural effusions, 2730
 in pleural mesothelioma
 preoperative evaluation of, 1951
 and radiation therapy, 1955-1956
 in radiation therapy, 2894-2897
 bronchiolitis obliterans in, 2896-2897
 clinical features in, 2895
 computed tomography in, 667, 2896
 diagnosis of, 2896
 diagnostic imaging in, 2895-2896
 dose-response curve in, 2894
 fibrosis in, 2895, 2896
 in lung cancer, 946
 genetic factors in, 2894
 histopathology in, 2895
 in Hodgkin's disease, 2377
 in lung cancer, 946, 995
 prevention of, 2894
 mechanism of lung injury in, 2894
 in pleural mesothelioma, 1955-1956
 pneumonitis in. *See* Pneumonitis, radiation-induced
 pulmonary function tests in, 2896
 time course of, 2895
 treatment of, 2896
 in stem cell transplantation, 2783-2784
 in tobacco smoke exposure, 552
 in transfusion-related acute lung injury, 2761
 Respiratory flora alterations, 2820
 Respiratory motions in radiation therapy for lung cancer, 947
 Respiratory syncytial virus, 2834t, 2837, 2853
 Response elements, 7
 Restriction enzymes, 5
 Rests, nephrogenic, intralobar and perilobar, 2172
 Resveratrol in cancer prevention, 592, 593t
 RET, 18, 202, 1730-1732
 in Hirschsprung's disease, 1731-1732
 in multiple endocrine neoplasia, 259, 617-618, 1730-1732, 1836, 1837
 locations of germline mutations in, 1731f
 in thyroid cancer, 18, 202, 259, 617-618, 1730, 1731t
 locations of germline mutations in, 1731f
 missense germline mutations in, 1730, 1731t
 in papillary carcinoma, 1734-1735, 1735f
 prophylactic thyroidectomy in, 1733, 1759
 testing for, 1759, 1837
 in children, presymptomatic, 3055
 Reticuloendotheliosis, leukemic, 2459
 Reticulosis, pagetoid, 2317, 2318
 Reticulum cell sarcoma
 of bone, 1929
 of central nervous system, 2330
 of skin, 2316
 Retinal pigment epitheliopathy, tumor-associated, in uveal melanoma, 2075
 Retinoblastoma, 1842, 2162-2163, 2182-2185
 associated anomalies in, 2182
 bilateral, 2183, 2184-2185
 limited disease, 2184
 chemotherapy in, 2185
 in children, 2162-2163, 2182-2185
 clinical presentation in, 2182-2183
 epidemiology of, 2169, 2170t, 2182
 etiology of, 2169-2170, 2170t
 exophytum and endophytum types, 2182
 genetic counseling in, 2182
 hereditary syndrome, 2098, 2162, 2182
 molecular genetic markers in, 653t
 risk assessment in, 652
 parental preconception exposures in, 2170t
 pathology in, 2182
 radiation therapy in, 2184-2185
 radiography in, 2183
 and *RB*, 2162-2163, 2169-2170, 2182.
 See also RB (Rb)
 second cancer risk in, 200, 2957, 2958f
 radiation-induced, 200, 2957
 sporadic, 2162, 2182
 staging of, 2183, 2183t, 2184t
 surgery in, 2184, 2185
 tumor suppressor genes in, 208-209, 2162, 2182
 unilateral, 2182, 2183
 advanced disease, 2184
 limited disease, 2184
cis-Retinoic acid
 in cerebral astrocytoma, 2126, 2127t
 in kidney cancer and interferon therapy, 1375, 1375t
 9-*cis*-Retinoic acid, 489-490, 576
 adverse effects of, 490
 in Kaposi's sarcoma, 490, 2578, 2581
 in lung cancer prevention, 584
 13-*cis*-Retinoic acid, 489, 490, 576
 adverse effects of, 490
 in esophageal cancer, 1067, 1067t
 in neuroblastoma, 2181
 in prevention of cancer, 489
 of head and neck, 489, 582, 818, 819
 of lung, 583, 943
 of skin, 584
 translational studies on, 587
 structure of, 576f
 all-*trans*-Retinoic acid (ATRA), 489, 576
 adverse effects of, 490-491, 2422
 cardiac toxicity in, 2909
 pulmonary toxicity in, 2901
 in cervical cancer prevention, 585
 in Kaposi's sarcoma, 2581
 in lung cancer, 920
 in myelogenous leukemia of children, 2243-2244
 in promyelocytic leukemia, acute, 21, 489, 2397-2398, 2422-2423
 pharmacology of, 489
 toxicity of, 491, 2422, 2901, 2909
 in skin cancer prevention, 584
 Retinoic acid receptors, 489
 RAR- α , 576
 in leukemia, 21, 489, 491, 2390t, 2397-2398
 RAR- β , 576
 in cervical cancer, 1523
 in lung cancer, 922
 in oral premalignancy, 586, 587, 587f, 791
 RAR- γ , 576, 587
 Retinoic acid syndrome in acute promyelocytic leukemia, 491, 2422
 Retinoid response elements, 489
 Retinoid X receptors, 489
 Retinoids, 489-491
 in cutaneous T-cell lymphoma, 2322, 2326-2327
 in interferon- α therapy, 467
 pharmacology of, 489-491, 576-577
 in prevention of cancer, 489, 575-588
 apoptosis-inducing activity in, 587
 biology and pharmacology of, 576-577
 of bladder, 585
 of breast, 584
 of cervix, 585
 clinical trials on, 577-587, 578t-579t
 of esophagus and stomach, 585
 of head and neck, 581-582, 791, 817-819
 translational studies on, 586-587




FIGURE 2-1. Molecular progression of cancer. Microscopic premalignant lesions originate within a background of hereditary predisposition. Genetic instability leads to molecular derangements that drive somatic genetic progression. Host interactions enable the cancer lesions to expand, vascularize, and metastasize. Diagnosis usually takes place at a late stage when there is a high probability of metastasis.

Separate genetic changes (beyond those causing uncontrolled growth) are required for tumor invasion and metastasis. Invasion and metastasis form a multistep cascade involving positive and negative regulatory pathways. Cancer invasion and angiogenesis are an uncontrolled version of physiologic invasion.

Genetic instability may predispose the premalignant cell to generate malignant offspring. Instability can take place at the macro level (chromosome karyotype), as well as the micro level (DNA sequence copy fidelity repair). Chromosomal rearrangement can activate silent oncogenes or delete regions containing suppressor genes. Loss of heterozygosity is a hallmark of suppressor gene inactivation in cancer progression. Telomerase defects may affect growth control as well as genetic instability. Mutations in cellular DNA can activate oncogenes or inactivate suppressor genes. Defects in DNA repair mechanisms contribute to the accumulation of genetic defects fueling cancer progression. Genetic defects causing an inhibition of cell death pathways are an important mechanism in tumorigenesis.

CANCER GENES: MODELS OF ACTION

Genetic alterations involved in cancer can activate inductive processes (oncogenes) or block negative pathways (suppressor genes). Early models of cancer genetics categorized cancer genes into oncogenes, which are growth inducing, and tumor suppressor genes, which are growth suppressing. Thus, mutations in oncogenes activate a promoting function, but lesions in tumor suppressors inactivate an inhibitory function. Exam-

ples of these models were the *ras* oncogene and the retinoblastoma (*rb*) tumor suppressor gene. Mutations in codons 12, 13, and 61 in the *ras* gene result in biochemical activation of the protein product and an induction of its transforming activity. Deletions or inactivating mutations in the *rb* gene lead to a compromised suppressor protein that is incapable of inhibiting cell growth. Aberrations in both genes are found as somatic mutations in human cancers and, in the case of *rb*, also in the germline of individuals at risk for cancer.

Dominant oncogenes play a significant role in human cancers. *Ras* mutations are found in 10% of cancers and appear frequently in colon and lung adenocarcinomas. *Ret* is a receptor tyrosine kinase in which activating single nucleotide mutations are associated with hereditary thyroid carcinomas. *Myc*, encoding a nuclear oncoprotein, is involved in the t(8;14) translocation, which is etiologic for Burkitt's lymphoma. Inappropriate overexpression of *myc* is sufficient for transformation of lymphocytes in transgenic models. Similarly, amplification and overexpression of the *HER2/neu* receptor tyrosine kinase not only causes mammary malignancies, but is prognostic in human breast cancers. Although originally these oncogene abnormalities were thought to induce cancer primarily through unregulated growth, other cellular phenotypes such as enhanced survival and motility may be equally important contributors to the cancer state.

It was also originally thought that tumor suppressor genes function mainly by inhibiting cell growth. Later, this was expanded to genes that block the emergence of a tumor, but not growth in culture. More recent studies, however, have uncovered other mechanisms unrelated to growth by which *tumor suppressor*



FIGURE 38.1-3. Photomicrograph of medullary thyroid carcinoma showing nests and sheets of small, uniform cells with scant to moderate amounts of amphophilic cytoplasm infiltrating around normal thyroid follicles. (Reprinted from Moley JF, Lairmore TC, Phay JE, Hereditary endocrinopathies. *Curr Probl Surg* 1999;36:653, with permission.)

In MEN 2B, 40% to 50% of patients develop pheochromocytomas, and all individuals develop neural gangliomas, particularly in the mucosa of the digestive tract, conjunctiva, lips, and tongue. MEN 2B patients also have megacolon, skeletal abnormalities, and markedly enlarged peripheral nerves. MEN 2B patients do not develop HPT. MTC develops at a very young age (infancy) and appears to be the most aggressive form of hereditary MTC, although its aggressiveness may be related more to the extremely early age of onset than to the biologic virulence of the tumor. MTC in MEN 2B is rarely curable.

FMTC is characterized by the development of MTC without any other endocrinopathies.²⁶ MTC in these patients has a later age of onset and follows a more indolent clinical course than MTC in patients with MEN 2A and MEN 2B. Occasional patients with FMTC never manifest clinical evidence of MTC (symptoms or a lump in the neck), though biochemical testing and histologic evaluation of the thyroid usually demonstrates MTC.

Genetics

THE RET PROTOONCOGENE. In 1987, the gene for MEN 2A was localized to the pericentromeric region of chromosome

10 (10q11.2) by linkage analysis.^{27,28} Subsequent studies demonstrated that the predisposition gene for MEN 2B and FMTC mapped to the same region as MEN 2A.^{29,30} The *RET* protooncogene resides within this critical region, which made it an obvious candidate gene for the MEN 2 syndromes.

The *RET* protooncogene was first discovered based on its ability to transform mouse NIH 3T3 fibroblasts in culture.³¹ The transforming *RET* sequences first identified represented a rearrangement of *RET* that occurred *in vitro* during the transfection assay.³² Sequence analysis of the *RET* protooncogene showed that it is a member of the receptor tyrosine kinase gene family.³³ *RET* was mapped to the proximal region of the long arm of chromosome 10 in 1989³⁴ and was shown to be expressed at high levels in both MTCs and pheochromocytomas.³⁵

RET mutations were identified in the constitutional DNA of MEN 2A and FMTC patients in 1993.^{36,37} Mulligan et al.³⁶ identified *RET* mutations in association with MEN 2 using an expression-based mutational analysis system. Analysis of genomic DNA from MEN 2A family members proved that the variants identified in the complementary DNAs (cDNAs) were also in the constitutional DNA of affected family members and not in individuals unaffected by MEN 2A. All mutations resulted in substitution of cysteine residues clustered near the transmembrane domain of *RET*. In all but one, MEN 2A tumor heterozygosity for the mutant and wild-type *RET* allele was retained. Mulligan et al.³⁶ suggested a dominant or dominant-negative mechanism for *RET* mutation in the development of MTC and pheochromocytomas in MEN 2A.³⁶

Donis-Keller et al.³⁷ used single-strand conformational variant (SSCV) and sequence analysis to identify mutations in *RET* in the constitutional DNA of patients with MEN 2A and FMTC. Both MEN 2A and FMTC were shown to be associated with mutations that result in substitution of cysteine residues in the extracellular portion of *RET* immediately adjacent to the transmembrane domain. Unexpectedly, the same mutations were found to characterize MEN 2A and FMTC kindreds.

Subsequent investigations demonstrated mutations of *RET* in MEN 2B patients.^{38,39} In 95% of cases of MEN 2B, the *RET* protooncogene mutation is a missense methionine-to-threonine (ATG to ACG) change at codon 918 in exon 16. This codon is positioned within the tyrosine kinase catalytic core of the intracellular domain. Two families with MEN 2B have been described that have a codon 883 mutation in the tyrosine kinase domain of *RET*.⁴⁰

More than 30 missense mutations have been described in MEN 2A and FMTC kindreds (Table 38.1-2).^{41,42} Most of these mutations result in nonconservative changes in cysteine residues, although changes in Glu, Val, Met, Leu, and Tyr have also been described. Several of these mutations have been shown to result in "gain-of-function" in the *RET* protein product, with increased intrinsic tyrosine kinase activity or alterations of substrate recognition (or both) and transforming capability.⁴³ The *RET* protooncogene encodes a protein with three domains: a cysteine-rich extracellular receptor domain, a hydrophobic transmembrane domain, and an intracellular tyrosine kinase catalytic domain (Fig. 38.1-4). The *RET* gene consists of at least 20 exons⁴⁴ and is expressed as five major mRNA species.^{45,46} The *RET* gene product is expressed in a limited number of cell types in the normal individual, including the thyroid C cells, the adrenal medulla, and parts of the brain. The gene is important in the embryonic development of the enteric nervous system and the kidneys.⁴⁷

TABLE 38.1-2. *RET* Mutations in Hereditary Medullary Thyroid Carcinoma

Syndrome	Missense Germline Mutations in the <i>RET</i> Protooncogene	
	Exon	Codon
MEN 2A, FMTC	10	609
		611
		618
		620
	11	631 ^a
		634
	13	790
FMTC	11	791
		630
		768
		804
	14	844 ^a
MEN 2B	15	891
	16	918
		883

FMTC, familial, non-MEN medullary thyroid carcinoma; MEN 2A, multiple endocrine neoplasia type 2A; MEN 2B, multiple endocrine neoplasia type 2B.

^aClinical features not yet characterized.

(Reprinted from Moley JF, Lairmore TC, Phay JE. Hereditary endocrinopathies. *Curr Probl Surg* 1999;36:653, with permission.)

Glial-derived neurotrophic factor (GDNF) and neurturin are ligands to the receptor domain of the *RET* gene product.⁴⁸⁻⁵⁴ GDNF is a 32-kD protein dimer that was first purified from glial cell lines and is a potent neurotrophic survival factor for motor neurons. A glycosylphosphatidylinositol-linked protein called *GDNF receptor-α* (GDNFR- α) is a cofactor in the signaling het-

erodimeric complex with *RET*. Current evidence suggests that GDNF binds directly to GDNFR- α and indirectly with *RET*. When triggered by ligand, wild-type *RET* dimerizes with another *RET* molecule, and this dimerization is responsible for phosphorylation and activation of the tyrosine kinase domain, with subsequent downstream signal transduction events. *RET* molecules that contain MEN 2A-type mutations are constitutively dimerized and therefore activated. In contrast, the mutation responsible for MEN 2B does not result in constitutive dimerization but changes the substrate specificity of the tyrosine kinase domain, which results in transformation.

OTHER *RET* GENOTYPE-PHENOTYPE CORRELATIONS

Cutaneous Lichen Amyloidosis. Interscapular lesions of cutaneous lichen amyloidosis (CLA) have been described in several families with MEN 2A.²² The total number of patients described with this entity is fewer than 100. In a 634→Tyr mutation was reported in one family with MEN 2A and CLA, and in two other families, a 634→Arg mutation was described that segregated with both *MEN2A* and CLA.⁵⁵

Hyperparathyroidism. HPT in MEN 2A is caused by parathyroid hyperplasia; the hypercalcemia is mild and often asymptomatic. HPT clusters in some families with MEN 2A. Whether specific *MEN2A* mutations are associated with a higher incidence of HPT remains controversial. Mulligan et al.⁵⁶ previously described a strong correlation between C634R mutation and HPT in families with MEN 2A, but other studies have been unable to confirm this relationship definitively.⁵⁷⁻⁵⁹

Hirschsprung's Disease. Hirschsprung's disease is characterized by absence of autonomic ganglion cells within the distal colonic parasympathetic plexus, resulting in obstruction and proximal megacolon. Approximately 80% of Hirschsprung's disease cases are sporadic, and the remainder are familial. A subset of familial Hirschsprung's cases have been found to be associated with germline mutations of *RET*.^{24,56} Most of these are inactivating and loss-




FIGURE 38.1-4. Diagram of *RET* gene product delineating locations of germline mutations found in multiple endocrine neoplasia type 2A and familial, non-MEN medullary thyroid carcinoma (ovals), germline mutations in multiple endocrine neoplasia type 2B (diamond), and mutations in hereditary Hirschsprung's disease (triangles). As mentioned in the text, the *RET* gene product is thought to create a dimer that forms a complex with glial-derived neurotrophic factor receptor- α or neurturin. The *RET* gene product is divided into the intracellular, transmembrane, and extracellular domains. ATP, adenosine triphosphate. (Adapted from Moley JF, Kim S, *Molecular genetics in surgical oncology*. Austin: RG Landes, 1994, with permission.)

TABLE 38.1-3. Loss of Heterozygosity in Pheochromocytomas and Medullary Thyroid Carcinomas

Study	Chromosomal Arms Tested	LOH in Pheos (No. LOH/No. Informative)	LOH in MTCs (No. LOH/No. Informative)
Khosla et al. ¹⁷⁸	1p, 2p, 3p, 5q, 10q, 13q, 16p, 17p, 17q, 22q	1p-13/31, 2p-1/34, 3p-4/24, 10q-1/22, 17p-7/27, 22q-5/18	1p-1/11, 22-1/7
Moley et al. ⁶⁹	1p, 1q	1p-12/18 (9/9 from MEN 2A and 2B patients)	1p-3/24
Yang et al. ¹⁷⁹	1p	1p-5/8	
Mathew et al. ¹⁸⁰	1p, 1q, 5, 6, 7, 11, 12	1p-4/6	1p-3/8
Shin et al. ¹⁸¹	1p, 22q	1p-12/22, 22q-8/20	
Mulligan et al. ⁷⁰	1-22, both arms	1p-15/25, 3p-10/18, 3q-9/15, 5q-1/7, 6q-1/7, 8p-1/8, 11p-3/19, 11q-2/23, 13-2/17, 17p-3/20, 17q-1/20, 22-8/20	1p-7/28, 3p-1/19, 3q-2/14, 7p-1/17, 10p-1/18, 10q-1/25, 11p-1/16, 13-2/22, 15-1/9, 21-1/8, 22-4/22
Dou et al. ⁷³	2-23	3q-13/13 F, 6/8 S, 21q-4/6 F, 2/7 S, 22q-7/13 F, 1/10 S, 11q-2/6 F, 3/7 S	3q-2/7 F, 1/8 S, 22q-4/15 F, 2/8 S
Herfarth et al. ⁷⁴	17p	17p-4/22	17p-0/14

F, familial (MEN 2A and 2B); LOH, loss of heterozygosity; MEN 2A, multiple endocrine neoplasia type 2A; MEN 2B, multiple endocrine neoplasia type 2B; MTCs, medullary thyroid carcinomas; Pheos, pheochromocytomas; S, sporadic (MEN 2A and 2B).

Note: If arm tested is not listed in LOH column, LOH was not found.

(Adapted from J Moley, Molecular events in the development and progression of medullary thyroid cancer and pheochromocytoma. In: Nelkin B, ed., *Genetic mechanisms in multiple endocrine neoplasia type 2*. Austin: RG Landes, 1996, with permission.)

of-function (frameshift and nonsense) mutations and are not associated with the MEN 2A phenotype. Several families, however, have been described in which Hirschsprung's disease cosegregates with either *MEN2A* or *FMTC* (missense codon 618 or 620 mutations).⁶⁰ Additionally, a few Hirschsprung's disease patients have been described with missense mutations in codon 609 or 620 who have no evidence of MEN 2A or MTC.^{23,25,56} It is interesting to note that the Hirschsprung's disease phenotype can be associated with either loss-of-function or gain-of-function mutations of *RET*. All patients with MEN 2B (missense codon 918 mutation) have megacolon and chronic colonic motility disturbances, though they usually do not require surgery for this (see Fig. 38.1-2C).⁶¹

RET Mutations in Sporadic Tumors. Mutations in the *RET* protooncogene have also been found in sporadic MTCs.^{37,40,62-64} The most frequent mutation in sporadic MTCs is the M918T mutation found in MEN 2B. Mutations have been found in other regions of the extracellular and intracellular domains. Missense, deletions, and insertion mutations have been described.^{41,65} Somatic *RET* mutations in sporadic pheochromocytomas are unusual but have also been described.⁴¹

Other Dominant Oncogenes in Medullary Thyroid Carcinomas and Pheochromocytomas. Absence of amplification of *N-MYC*, *C-MYC*, and *ERB-B2* has been reported in MTCs and pheochromocytomas. Roncalli et al.⁶⁶ reported that *N-myc* expression in more than 10% of tumor cells, as detected by immunohistochemistry, was associated with poorer survival, sporadic disease, and male gender. These investigators found no evidence of gene amplification and did not determine the basis for the overexpression.⁶⁶ Our group reported absence of mutation of the *H-RAS*, *N-RAS*, and *K-RAS* genes in a series of pheochromocytomas and MTCs analyzed by direct sequencing.⁶⁷ Likewise, examination of nerve growth factor and nerve growth factor receptor (p75) showed no abnormality at the DNA or RNA levels.⁶⁸

Other Tumor Suppressor Genes in Medullary Thyroid Carcinoma and Pheochromocytoma. Several studies have evaluated loss of heterozygosity (LOH) at tumor suppressor loci in pheochromocytomas

and MTCs; these are summarized in Table 38.1-3. The cumulative data indicate a higher-than-background incidence of LOH in pheochromocytomas on chromosome arms 1p, 3p, 17p, and 22q.^{69,70} In MTCs, the report by Mulligan et al.⁷⁰ suggests a significant incidence of 1p LOH; however, evaluation of other chromosomal arms yielded no consistent findings. Lack of significant LOH on 10q, at the *RET* locus, supports the hypothesis that the *RET* protooncogene acts as a dominant oncogene as opposed to a tumor suppressor gene.⁷¹ Chromosome 1 is the largest chromosome, and LOH analysis on 1p in pheochromocytomas suggests a very large region of deletion. Our studies have indicated that the entire short arm of 1p is lost in pheochromocytomas from patients with MEN 2A and MEN 2B.⁶⁹ Fine mapping of the region of deletion suggests a possible common breakpoint in the centromeric region defined by the markers D1S514 and D1S442.⁷²

The high rate of LOH on 3p in pheochromocytomas suggests an as-yet undefined tumor suppressor locus.^{70,73} LOH on 17p suggests possible involvement of the *TP53* gene.⁷⁴ Existing reports on *TP53* mutations in pheochromocytomas are conflicting. Two Japanese groups reported no evidence of *TP53* mutations in pheochromocytomas.^{75,76} In contrast, a Chinese group reported *TP53* mutations in five of six tumors tested.⁷⁷ Four of these mutations were in exon 4. Our group reported a series of 22 pheochromocytomas and 29 MTCs that were screened with four different markers for LOH on 17p.⁷⁴ SSCV analysis of exons 4 through 9 of the *TP53* gene was performed in 20 of the pheochromocytomas and in 22 of the MTCs. The expression of p53 was determined by immunohistochemistry in 19 pheochromocytomas and in 17 MTCs, using two different antibodies (D01 and D07) on both frozen and paraffin-embedded tissues. LOH was demonstrated on 17p in 4 of the 22 pheochromocytomas and in none of the MTCs. No mutations were detected in any of the tumors screened by SSCV analysis. Immunohistochemical staining of frozen and paraffin-embedded tumor sections did not show p53 overexpression in any of the tumors examined. These findings indicate that mutations in the *TP53* gene are an uncommon event in the tumorigenesis of hereditary and sporadic pheochromocytomas and MTCs.⁷⁴

Pheochromocytomas also occur in neurofibromatosis type 1 (NF1) and von Hippel-Lindau (VHL) disease, both of which

CONTRIBUTING AUTHORS

James L. Abbruzzese, MD

Annie Laurie Howard Research Distinguished Professor
Department of Gastrointestinal Medical Oncology
The University of Texas M. D. Anderson Cancer Center
Houston, Texas

Gregory P. Adams, PhD

Assistant Member
Department of Medical Oncology
Fox Chase Cancer Center
Philadelphia, Pennsylvania

Daniel M. Albert, MD, MS

F. A. Davis Professor and Chair; Lorenz E. Zimmerman Professor
Department of Ophthalmology and Visual Sciences
University of Wisconsin Center for Health Sciences
Madison, Wisconsin

Kaled M. Alektiar, MD

Assistant Attending
Department of Radiation Oncology
Memorial Sloan-Kettering Cancer Center
New York, New York

H. Richard Alexander, MD

Head, Surgical Metabolism Section
Surgery Branch
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Carmen J. Allegra, MD

Chief, Medicine Branch
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Nasser K. Altorki, BCh, MB

Professor of Cardiothoracic Surgery
Weill Medical College of Cornell University
New York, New York

Howard Ira Amols, PhD

Professor of Physics in Radiology
Weill Medical College of Cornell University
Memorial Sloan-Kettering Cancer Center
New York, New York

Susan Anderson, BSN, MFA

Director, Clinical Trials Office
Yale Cancer Center
Yale University School of Medicine
New Haven, Connecticut

Karen H. Antman, MD

Wu Professor of Medicine and Pharmacology
Herbert Irving Comprehensive Cancer Center
Columbia University College of Physicians and Surgeons
New York, New York

James O. Armitage, MD

Professor of Medicine
Dean of the College of Medicine
University of Nebraska Medical Center
Omaha, Nebraska

Susanne M. Arnold, MD

Assistant Professor of Medicine
Department of Hematology and Oncology
Lucille P. Markey Cancer Center
University of Kentucky Chandler Medical Center
Lexington, Kentucky

Michael B. Atkins, MD

Associate Professor of Medicine
Department of Hematology/Oncology
Harvard Medical School
Beth Israel Deaconess Medical Center
Boston, Massachusetts

Richard M. Auchter, MD

Clinical Assistant Professor of Human Oncology
University of Wisconsin Medical School
Madison, Wisconsin

David J. Austin, PhD

Assistant Professor of Chemistry
Yale University School of Medicine
New Haven, Connecticut

Dean F. Bajorin, MD

Professor of Medicine
Weill Medical College of Cornell University
Attending Physician
Memorial Sloan-Kettering Cancer Center
New York, New York

Allen E. Bale, MD

Associate Professor of Genetics
Yale University School of Medicine
New Haven, Connecticut

Bart Barlogie, MD, PhD

Professor of Medicine and Pathology
Director, Arkansas Cancer Research Center
University of Arkansas for Medical Sciences
Little Rock, Arkansas

David L. Bartlett, MD

Surgery Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Susan E. Bates, MD

Head, Molecular Therapeutics Section, Medicine Branch
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Jonathan S. Berek, MD

Professor and Chair of Applied Anatomy
Vice Chair and Chief, Division of
Gynecologic Oncology
University of California, Los Angeles, UCLA School
of Medicine
Los Angeles, California

Ann M. Berger, RN, MSN, MD

Chief of Pain and Palliative Care Service
National Institutes of Health
Bethesda, Maryland

Philip Bierman, MD

Associate Professor of Medicine
Department of Internal Medicine
Section of Oncology/Hematology
University of Nebraska Medical Center
Omaha, Nebraska

Peter McLaren Black, MD, PhD

Franc D. Ingraham Professor of Neurosurgery
Harvard Medical School
Brigham and Women's Hospital
Children's Hospital
Boston, Massachusetts

Clara D. Bloomfield, MD

Professor of Medicine
Ohio State University College of Medicine
and Public Health
Department of Internal Medicine
Division of Hematology/Oncology
The Comprehensive Cancer Center
Columbus, Ohio

David A. Bluemke, MD, PhD, MSB

Associate Professor of Radiology
Johns Hopkins University School of Medicine
Baltimore, Maryland

George J. Bosl, MD

Chair, Department of Medicine
Memorial Sloan-Kettering Cancer Center
New York, New York

Douglas E. Brash, PhD

Professor of Therapeutic Radiology and Genetics
Yale University School of Medicine
New Haven, Connecticut

Murray F. Brennan, MD

Chair, Department of Surgery
Memorial Sloan-Kettering Cancer Center
New York, New York

Holly K. Brown, MD

Instructor of Orthopaedic Surgery
Memorial Sloan-Kettering Cancer Center
New York, New York

Suzanne J. Brown, MD

Fellow in Oncology
Yale Comprehensive Cancer Center
Yale University School of Medicine
New Haven, Connecticut

Nick Bryan, MD, PhD

Professor of Radiology
University of Pennsylvania School of Medicine
Philadelphia, Pennsylvania

Gary L. Buchsacher, Jr., MD, PhD

Department of Medicine
Division of Hematology/Oncology
University of California, San Diego, School
of Medicine
La Jolla, California

Thomas W. Burke, MD

Professor of Gynecologic Oncology
The University of Texas M. D. Anderson
Cancer Center
Houston, Texas

Chandra Burman, PhD

Associate Attending Physician
Department of Medical Physics
Memorial Sloan-Kettering Cancer Center
New York, New York

Michael A. Caligiuri, MD

Professor of Medicine
Ohio State University College of Medicine
and Public Health
Arthur James Cancer Hospital and Research Institute
Columbus, Ohio

Robert B. Cameron, MD

Assistant Professor of Surgery
University of California, Los Angeles, UCLA School
of Medicine
Los Angeles, California

Lewis C. Cantley, PhD

Professor of Cell Biology
Harvard Medical School
Beth Israel Deaconess Medical Center
Boston, Massachusetts

Michele Carbone, MD, PhD

Associate Professor of Pathology
Department of Pathology/Cancer Immunology
Loyola University Medical Center
Maywood, Illinois

Christopher L. Carpenter, MD, PhD

Assistant Professor of Medicine
Harvard Medical School
Beth Israel Deaconess Medical Center
Boston, Massachusetts

Peter R. Carroll, MD

Professor and Chair, Department of Urology
University of California, San Francisco, School
of Medicine
San Francisco, California

John A. Carucci, MD, PhD

Clinical Instructor of Dermatology
Yale University School of Medicine
New Haven, Connecticut

Webster K. Cavenee, PhD

Professor of Medicine
University of California, San Diego, School of Medicine
Director, Ludwig Institute for Cancer Research
Center for Molecular Genetics; University of California
Laboratory of Tumor Biology
La Jolla, California

Bruce A. Chabner, MD

Professor of Medicine
Harvard Medical School
Department of Hematology/Oncology
Massachusetts General Hospital
Boston, Massachusetts

R. S. K. Chaganti, PhD

Member and William E. Snee Chair
Human Genetics
Memorial Sloan-Kettering Cancer Center
New York, New York

Setsuko Kuki Chambers, MD

Professor of Obstetrics and Gynecology
Division of Gynecologic Oncology
Yale University School of Medicine
New Haven, Connecticut

Lisa Chertkov, MD

Fellow in Psychiatry and Behavioral Sciences
Memorial Sloan-Kettering Cancer Center
New York, New York

Bruce D. Cheson, MD

Clinical Professor of Medicine
Georgetown University School of Medicine
Vincent T. Lombardi Cancer Research Center
Washington, DC
Head, Medicine Section
Cancer Therapy Evaluation Program
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Richard W. Childs, MD

Senior Investigator
Hematology Branch
National Heart, Lung, and Blood Institute,
National Institutes of Health
Bethesda, Maryland

Edward Chu, MD

Associate Professor of Medicine and Pharmacology
Yale Cancer Center
Yale University School of Medicine
Director, Veterans Administration Connecticut
Cancer Center
New Haven, Connecticut

Chen Chui, PhD

Associate Professor of Medical Physics
Memorial Sloan-Kettering Cancer Center
New York, New York

Rebecca A. Clark-Snow, RN, BSN, OCN

Director of Clinical Services
Oncology and Hematology Associates
of Kansas City
Kansas City, Missouri

Daniel G. Colt, MD

Chief of Gastric and Mixed Tumor Service
Department of Surgery
Memorial Sloan-Kettering Cancer Center
New York, New York

Graham A. Colditz, MD, DrPH

Professor of Medicine
Harvard Medical School
Brigham and Women's Hospital
Boston, Massachusetts

Phillip Cole, MD, PhD

Professor Emeritus
Department of Epidemiology
University of Alabama School of Medicine
Birmingham, Alabama

Oliver Michael Colvin, MD

Professor of Hematology/Oncology
Director, Duke Comprehensive Cancer Center
Duke University Medical Center
Durham, North Carolina

Dennis L. Cooper, MD

Associate Professor of Medicine
Yale University School of Medicine
Department of Medical Oncology
Yale-New Haven Hospital
New Haven, Connecticut

Carlos Cordon-Cardo, MD, PhD

Member and Director, Division
of Molecular Pathology
Department of Pathology
Memorial Sloan-Kettering Cancer Center
New York, New York

José Costa, MD

Professor of Pathology
Yale University School of Medicine
Yale-New Haven Hospital
New Haven, Connecticut

Guido Dalbagni, MD, FACS

Assistant Professor of Urology
Weill Medical College of Cornell University
Memorial Sloan-Kettering Cancer Center
New York, New York

Riccardo Dalla-Favera, MD

Uris Professor of Pathology and Genetics
Director, Institute of Cancer Genetics
Columbia University College of Physicians
and Surgeons
New York, New York

Ramsey M. Dallal, MD

Resident of Surgery
University of Pittsburgh Medical Center
Pittsburgh, Pennsylvania

Christopher Daugherty, MD

Assistant Professor of Medicine
University of Chicago Pritzker School
of Medicine
Chicago, Illinois

Lisa M. DeAngelis, MD

Chair, Department of Neurology
Memorial Sloan-Kettering Cancer Center
New York, New York

Albert B. Deisseroth, MD, PhD

Ensign Professor of Medicine
Department of Internal Medicine
Yale University School of Medicine
New Haven, Connecticut

Vincent T. DeVita, Jr., MD

Director, Yale Cancer Center
Professor of Medicine
Professor of Epidemiology and Public Health
Yale University School of Medicine
New Haven, Connecticut

Robert B. Dickson, PhD

Professor of Oncology
Georgetown University School of Medicine
Vincent T. Lombardi Cancer Research Center
Washington, DC

Volker Diehl, MD PhD

Professor of Medicine
Department of Internal Medicine I
University Hospital Cologne
Köln, Germany

Joseph J. Disa, MD

Assistant Attending Surgeon
Department of Surgery/Plastic Surgery
Memorial Sloan-Kettering Cancer Center
New York, New York

Gerard M. Doherty, MD

Associate Professor of Surgery
Washington University School of Medicine
St. Louis, Missouri

Sarah S. Donaldson, MD, FACP

Professor of Radiation Oncology
Stanford University School of Medicine
Stanford, California

David H. Ebb, MD

Assistant Professor of Pediatrics
Harvard Medical School
MassGeneral Hospital for Children
Boston, Massachusetts

Richard L. Edelson, MD

Professor and Chair, Dermatology
Yale University School of Medicine
New Haven, Connecticut

Patricia J. Eifel, MD

Professor of Radiation Oncology
The University of Texas M. D. Anderson Cancer Center
Houston, Texas

Lee M. Ellis, MD

Associate Professor of Surgery and Cancer Biology
Department of Surgical Oncology and Cancer Biology
The University of Texas M. D. Anderson Cancer Center
Houston, Texas

Charles Erlichman, MD, FRCP(C)

Professor of Oncology
Department of Medical Oncology
Mayo Medical School
Mayo Clinic
Rochester, Minnesota

Elihu Estey, MD

Professor of Medicine
Department of Leukemia
The University of Texas M. D. Anderson Cancer Center
Houston, Texas

Douglas B. Evans, MD

Professor of Surgery
Department of Surgical Oncology
The University of Texas M. D. Anderson Cancer Center
Houston, Texas

Stefan Faderl, MD

Assistant Professor of Medicine
Department of Leukemia
The University of Texas M. D. Anderson Cancer Center
Houston, Texas

Eric R. Fearon, MD, PhD

Associate Professor of Internal Medicine, Human Genetics,
and Pathology
Department of Internal Medicine, Division of
Medical Genetics
University of Michigan Medical School
Ann Arbor, Michigan

Isaiah J. Fidler, DVM, PhD

Professor of Cancer Biology and Urology
The University of Texas M. D. Anderson
Cancer Center
Houston, Texas

Elliot K. Fishman, MD

Professor of Radiology and Oncology
Johns Hopkins University School of Medicine
Johns Hopkins Hospital
Baltimore, Maryland

Frederick A. Flatow, MD, FACP

Assistant Clinical Professor of Medicine
Department of Oncology
Yale University School of Medicine
New Haven, Connecticut
Medical Director, The Connecticut Hospice
Branford, Connecticut

John C. Flickinger, MD

Professor of Radiation Oncology and
Neurological Surgery
Department of Radiation Oncology
University of Pittsburgh Medical Center—
Presbyterian Hospital
Pittsburgh, Pennsylvania

Miklos C. Fogarasi, MD

Senior Fellow in Medicine, Section of Medical Oncology
Yale University School of Medicine
Yale Cancer Center
New Haven, Connecticut

Kathleen M. Foley, MD

Professor of Neurology, Neuroscience, and
Clinical Pharmacology
Weill Medical College of Cornell University
Attending Neurologist
Memorial Sloan-Kettering Cancer Center
New York, New York

Judah Folkman, MD

Andrus Professor of Pediatric Surgery
Professor of Cell Biology
Harvard Medical School
Children's Hospital
Boston, Massachusetts

Yuman Fong, MD

Attending Surgeon
Memorial Hospital
Member, Memorial Sloan-Kettering Cancer Center
New York, New York

Kwun M. Fong, MD

Hamon Center for Therapeutic Oncology Research
University of Texas Southwestern Medical Center
at Dallas
Dallas, Texas

Kenneth A. Foon, MD

Professor of Medicine
Department of Internal Medicine
University of Cincinnati College of Medicine
Director, Barrett Cancer Center
Cincinnati, Ohio

Arlene A. Forastiere, MD

Professor of Oncology
Johns Hopkins University School of Medicine
Baltimore, Maryland

Douglas L. Fraker, MD

Jonathon Rhoads Associate Professor of Surgery
Chief, Division of Surgical Oncology
University of Pennsylvania School of Medicine
Philadelphia, Pennsylvania

Zvi Y. Fuks, MD

Department of Radiation Oncology
Memorial Sloan-Kettering Cancer Center
New York, New York

Brian G. Fuller, MD

Senior Investigator
Radiation Oncology Branch
National Cancer Institute, Radiation Oncology
Sciences Program
National Institutes of Health
Bethesda, Maryland

Gianluca Galdano, MD, PhD

Department of Medical Sciences
Division of Internal Medicine
Amedeo Avogadro University of Eastern Piedmont
Novara, Italy

Don Ganem, MD

Professor of Microbiology and Medicine
Howard Hughes Medical Institute
University of California, San Francisco, School
of Medicine
San Francisco, California

Patricia A. Ganz, MD

Professor, School of Medicine and Public Health
Department of Medicine (Hematology-Oncology)
and Health Services
University of California, Los Angeles, UCLA School of Medicine
University of California, Los Angeles, Jonsson
Comprehensive Cancer Center
Los Angeles, California

Alan C. Geller, MPH, RN

Research Assistant Professor of Dermatology
Boston University School of Medicine
Boston, Massachusetts

Lynn Gerber, MD

Clinical Professor of Medicine
Georgetown University School of Medicine
Chief, Rehabilitation Medicine Department
Clinical Center, National Institutes of Health
Bethesda, Maryland

Jean-Francois H. Geschwind, MD

Assistant Professor of Radiology and Surgery
Johns Hopkins University School of Medicine
Director, Interventional Radiology Research
Department of Radiology and Surgery
Johns Hopkins Hospital
Baltimore, Maryland

Robert J. Ginsberg, MD, FRCS(C)

Professor of Surgery
Weill Medical College of Cornell University
Memorial Sloan-Kettering Cancer Center
New York, New York

Eli Glatstein, MD

Professor of Radiation Oncology
University of Pennsylvania School of Medicine
Hospital of the University of Pennsylvania
Philadelphia, Pennsylvania

Frank A. Greco, MD

Director, Sarah Cannon Cancer Center
Centennial Medical Center
Nashville, Tennessee

Daniel M. Green, MD

Professor of Pediatrics
Roswell Park Cancer Institute
University at Buffalo School of Medicine and Biomedical Sciences
Buffalo, New York

Peter Greenwald, MD, DrPH

Director, Division of Cancer Prevention
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Michael R. Grever, MD

Professor of Internal Medicine and Hematology and Oncology
Chairman, Department of Internal Medicine
Program Leader for the Experimental Therapeutics Program
James Comprehensive Cancer Center
The Ohio State University
Columbus, Ohio

James D. Griffin, MD

Professor of Medicine
Harvard Medical School
Brigham and Women's Hospital
Department of Adult Oncology
Dana-Farber Cancer Institute
Boston, Massachusetts

Phillip H. Gutin, MD

Chief of Neurosurgery; Fred Lebow Chair of Neurooncology
Division of Surgery
Memorial Sloan-Kettering Cancer Center
New York, New York

Douglas Hageman, MPA

Chief, Office of Clinical Informatics
Division of Clinical Sciences
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

John D. Hainsworth, MD

Director, Clinical Research
Sarah Cannon Cancer Center
Centennial Medical Center
Nashville, Tennessee

Ulrike M. Hamper, MD, MBA

Associate Professor of Radiology and Oncology
Johns Hopkins University School of Medicine
Baltimore, Maryland

Jay R. Harris, MD

Professor of Radiation Oncology
Harvard Medical School
Chief of Radiation Oncology
Brigham and Women's Hospital,
Children's Hospital
Dana-Farber Cancer Institute
Boston, Massachusetts

Nancy Lee Harris, MD

Professor of Pathology
Massachusetts General Hospital
Boston, Massachusetts

Louis B. Harrison, MD

Professor of Radiation Oncology
Albert Einstein College of Medicine of
Yeshiva University
Chair, Radiation Oncology
Beth Israel Medical Center
St. Luke's-Roosevelt Hospital Center
New York, New York

Peter W. Heald, MD

Associate Professor of Dermatology
Yale University School of Medicine
New Haven, Connecticut

John H. Healey, MD

Chief, Orthopaedic Surgery Service
Department of Surgery
Memorial Sloan-Kettering Cancer Center
New York, New York

John D. Heiss, MD

Staff Physician
Surgical Neurology Branch
National Institute of Neurological Disorders and Stroke
National Institutes of Health
Bethesda, Maryland

Samuel Hellman, MD

A. N. Pritzker Distinguished Service Professor
Department of Radiation and Cellular Oncology
University of Chicago
Chicago, Illinois

Lee J. Helman, MD

Chief
Pediatric Oncology Branch
Division of Clinical Sciences
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Charles H. Hennekens, MD, DrPH

Visiting Professor
Department of Medicine and Epidemiology
and Public Health
University of Miami School of Medicine
Miami, Florida

Meenhard Herlyn, DVM, DSc
The Wistar Institute
Philadelphia, Pennsylvania

Harry W. Herr, MD
Attending Surgeon
Department of Urology
Weill Medical College of Cornell University
Memorial Sloan-Kettering Cancer Center
New York, New York

Robert A. Hlatt, MD, PhD
Deputy Director, Division of Cancer Control
and Population Sciences
National Cancer Institute, National Institutes
of Health
Rockville, Maryland

Jeanne Hicks, MD
Associate Professor of Internal Medicine
George Washington University School of Medicine
and Health Sciences
Associate Professor of Rehabilitation Medicine
Department of Orthopedic Surgery
Georgetown University School of Medicine
Washington, DC
Deputy Chief, Department of
Rehabilitation Medicines
National Institutes of Health
Bethesda, Maryland

Paulo M. Hoff, MD
Assistant Professor
Department of Gastrointestinal Medical Oncology
The University of Texas M. D. Anderson Cancer Center
Houston, Texas

John P. Hoffman, MD
Professor of Surgery
Temple University School of Medicine
Senior Member, Fox Chase Cancer Center
Philadelphia, Pennsylvania

Steven M. Holland, MD
Senior Clinical Investigator, Laboratory
of Host Defenses
National Institute for Allergies and Infectious Diseases
and Warren Grant Magnuson Clinical Center
National Institutes of Health
Bethesda, Maryland

McDonald K. Horne III, MD
Senior Clinical Investigator, Clinical Pathology Department
Warren Grant Magnuson Clinical Center, National
Institutes of Health
Bethesda, Maryland

Peter M. Howley, MD
George Fabyan Professor of Comparative Pathology
Department of Pathology
Harvard Medical School
Boston, Massachusetts

Susan Molloy Hubbard, RN, MPA
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Margle Hunt, MS
Assistant Attending
Department of Medical Physics
Memorial Sloan-Kettering Cancer Center
New York, New York

Patrick Hwu, MD
Senior Investigator, Division of Clinical Sciences
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Bonnie A. Indeck, MSW
Clinical Instructor of Medicine
Department of Social Work/Oncology
Yale University School of Medicine
Yale-New Haven Hospital
New Haven, Connecticut

William Isaacs, PhD
Professor of Urology and Oncology
Johns Hopkins University School of Medicine
Baltimore, Maryland

Robert T. Jensen, MD
Chief, Digestive Diseases Branch
National Institute of Diabetes and Digestive
and Kidney Diseases, National Institutes of Health
Bethesda, Maryland

Steven W. Johnson, PhD
Research Assistant Professor of Pharmacology
University of Pennsylvania School of Medicine
Philadelphia, Pennsylvania

Glenn W. Jones, MD, FRCP, MSc
Associate Professor of Medicine
Department of Radiation Oncology
McMaster University Undergraduate Medical Programme
School of Medicine
Hamilton, Ontario, Canada

Barry M. Kacinski, MD, PhD
Professor of Therapeutic Radiology, Obstetrics/Gynecology,
and Dermatology
Yale University School of Medicine
New Haven, Connecticut

Robert J. Kaner, MD
Associate Professor of Clinical Medicine
Department of Medicine, Division of Pulmonary
and Critical Care Medicine
Weill Medical College of Cornell University
New York Presbyterian Hospital
New York, New York

Christine Kannler, MD, MPH
Medical Resident, Internal Medicine Program
Boston University School of Medicine
Boston, Massachusetts

Hagop M. Kantarjian, MD
Professor of Medicine
Chair, Department of Leukemia
The University of Texas M. D. Anderson
Cancer Center
Houston, Texas

Phillip W. Kantoff, MD

Associate Professor of Medicine
Harvard Medical School
Director, Lauk Center for Genitourinary Oncology,
Adult Oncology
Dana-Farber Cancer Institute
Boston, Massachusetts

Martin S. Karpeh, MD

Assistant Attending
Department of Surgery, Gastric and Mixed Tumor Service
Memorial Sloan-Kettering Cancer Center
New York, New York

Michael B. Kastan, MD, PhD

Member and Chair, Department of
Hematology/Oncology
St. Jude Children's Research Hospital
Memphis, Tennessee

David P. Kelsen, MD

Professor of Medicine
Weill Medical College of Cornell University
Chief, Gastrointestinal Oncology Service
Memorial Sloan-Kettering Cancer Center
New York, New York

Nancy Kemeny, MD

Professor of Medicine
Weill Medical College of Cornell University
Attending Physician
Department of Medicine, Solid Tumor Division,
Gastrointestinal Oncology Service
Memorial Sloan-Kettering Cancer Center
New York, New York

Robert S. Kerbel, PhD

Professor and Director
Department of Medical Biophysics
Division of Cancer Biology Research
University of Toronto Faculty of Medicine
Sunnybrook and Women's College Health Sciences Centre
Toronto, Ontario, Canada

Elliott Kleff, MD, PhD

Albee Professor
Department of Medicine and Microbiology and Molecular Genetics
Harvard Medical School
Brigham and Women's Hospital
Boston, Massachusetts

Thomas J. Kilroy, MD

Director of Dental Oncology
Porter Care Hospital
Denver, Colorado

E. Edmund Kim, MD, MS

Professor of Radiology and Medicine
Department of Nuclear Medicine and Diagnostic Radiology
The University of Texas M. D. Anderson Cancer Center
Houston, Texas

John M. Kirkwood, MD

Professor of Medicine, Division of Hematology/Oncology
University of Pittsburgh School of Medicine
Vice Chair for Clinical Research
Director Melanoma Center
University of Pittsburgh Cancer Institute
Pittsburgh, Pennsylvania

David E. Kleiner, Jr., MD, PhD

Director, Clinical Operations
Laboratory of Pathology
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

M. Tish Knobf, RN, PhD

Associate Professor
American Cancer Society Professor of Oncology Nursing
Adult Advanced Practice Program
Yale University School of Nursing
New Haven, Connecticut

Howard K. Koh, MD, MPH

Adjunct Professor
Boston University School of Medicine, School of Public Health
Commissioner of Public Health, Commonwealth
of Massachusetts
Boston, Massachusetts

Stanley J. Korsmeyer, MD

Sidney Farber Professor of Pathology
Professor of Medicine
Harvard Medical School
Investigator, Howard Hughes Medical Institute
Director, Program in Molecular Oncology
Department of Cancer Immunology and AIDS
Dana-Farber Cancer Institute
Boston, Massachusetts

Robert C. Kurtz, MD

Professor of Clinical Medicine
Department of Medicine
Weill Medical College of Cornell University
Attending Physician
Memorial Sloan-Kettering Cancer Center
New York, New York

Terry C. Lairmore, MD

Associate Professor of Endocrine and Oncologic Surgery
Washington University School of Medicine
St. Louis, Missouri

Theodore S. Lawrence, MD, PhD

Isadore Lampe Professor and Chair
Department of Radiation Oncology
University of Michigan Medical School
Ann Arbor, Michigan

Hop N. Le, MD

Fellow in Surgery
University of California, San Francisco, School of Medicine
San Francisco, California

Fredrick Leach, MD, PhD

Urologic Oncology Branch
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Keith L. Lee, MD

Research Fellow in Prostate Cancer
Department of Urology
University of California, San Francisco, School of Medicine
San Francisco, California

David J. Leffell, MD

Professor of Dermatology and Surgery
Yale University School of Medicine
New Haven, Connecticut

Alan T. Lefor, MD, MPH

Associate Professor of Clinical Surgery
University of California, Los Angeles, UCLA School
of Medicine
Director, Division of Surgical Oncology
Cedars-Sinai Medical Center
Los Angeles, California

Steven A. Leibel, MD

Chair, Department of Radiation Oncology
Memorial Sloan-Kettering Cancer Center
New York, New York

Victor A. Levin, MD

Professor of Neurooncology
The University of Texas M. D. Anderson Cancer Center
Houston, Texas

Arthur E. Li, MD

Post-Doctoral Fellow in Radiology
Johns Hopkins Hospital
Baltimore, Maryland

Peining Li, PhD

Assistant Director, Cytogenetics Laboratory
Department of Genetics
Yale University School of Medicine
New Haven, Connecticut

Steven K. Libutti, MD

Senior Investigator
Surgery Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

W. Marston Linehan, MD

Chief, Urologic Surgery
Urologic Oncology Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

C. Clifton Ling, PhD

Enid Haupt Chair
Department of Medical Physics
Memorial Sloan-Kettering Cancer Center
New York, New York

Michael P. Link, MD

Professor of Pediatrics
Chief, Division of Hematology, Oncology and
Bone Marrow Transplantation
Stanford University School of Medicine
Stanford, California

Lance A. Liotta, MD, PhD

Chief, Laboratory of Pathology
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Marc E. Lippman, MD

Professor of Oncology and Medicine
Georgetown University School of Medicine
Director, Vincent T. Lombardi Cancer Research Center
Georgetown University Hospital
Washington, DC

Scott M. Lippman, MD

Professor of Medicine and Cancer Prevention
Department of Thoracic/Head and Neck Medical
Oncology, Department of Clinical Cancer Prevention
The University of Texas M. D. Anderson Cancer Center
Houston, Texas

Larry I. Lipshultz, MD

Professor of Urology
Chief, Division of Male Reproductive Medicine
and Surgery
Baylor College of Medicine
Houston, Texas

Richard F. Little, MD, MPH

Senior Clinical Investigator
HIV and AIDS Malignancy Branch
Division of Clinical Sciences
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Mark S. Litwin, MD, MPH

Associate Professor of Urology and Health Services
University of California, Los Angeles, UCLA School
of Medicine
Los Angeles, California

Edison T. Liu, MD

Director, Division of Clinical Sciences
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Virginia Livolsi, MD

Professor of Pathology and Laboratory Medicine
University of Pennsylvania School of Medicine
Philadelphia, Pennsylvania

Jay S. Loeffler, MD

Director, Northeast Proton Therapy Center
Department of Radiation Oncology
Massachusetts General Hospital
Boston, Massachusetts

Patrick J. Loehrer, MD

Professor of Medicine
Section of Hematology Oncology
Indiana University School of Medicine
Indianapolis, Indiana

Anthony Lomax, MSc, PhD

Doctor of Radiation Medicine
Paul Scherrer Institut
Villigen, Switzerland

Scott Long, MD, PhD

Assistant Clinical Professor of Medicine
Yale University School of Medicine
New Haven, Connecticut
Physician, The Connecticut Hospice
Branford, Connecticut

Charles L. Loprinzi, MD

Professor and Chair
Department of Medical Oncology
Mayo Medical School
Mayo Clinic
Rochester, Minnesota

Thomas LoSasso, MD

Memorial Sloan-Kettering Cancer Center
New York, New York

Michael T. Lotze, MD

Professor of Surgery, Molecular Genetics,
and Biochemistry
University of Pittsburgh School of Medicine
Pittsburgh, Pennsylvania

David N. Louis, MD

Associate Professor of Pathology
Department of Pathology and
Neurosurgical Service
Harvard Medical School
Massachusetts General Hospital
Boston, Massachusetts

Andrew M. Lowy, MD

Chief of Surgical Oncology
Associate Medical Director
Barrett Cancer Center
University of Cincinnati Medical Center
Cincinnati, Ohio

Douglas R. Lowy, MD

Chief, Laboratory of Cellular Oncology
Division of Basic Sciences
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Gigas Mageras, PhD

Associate Attending Physician
Department of Medical Physics
Memorial Sloan-Kettering Cancer Center
New York, New York

Robert G. Maki, MD, PhD

Clinical Assistant Physician
Department of Medicine
Memorial Sloan-Kettering Cancer Center
New York, New York

Martin M. Malawer, MD, FACS

Professor of Orthopedic Surgery
George Washington University School of Medicine
and Health Sciences
Washington Hospital Center, Washington Cancer Institute
Washington, DC
Consultant, Surgery Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

David Malkin, MD

Associate Professor of Pediatrics
Division of Hematology/Oncology
University of Toronto Faculty of Medicine
The Hospital for Sick Children
Toronto, Ontario, Canada

Francesco M. Marincola, MD

Senior Investigator
Surgery Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Peter Maslak, MD

Assistant Professor
Medicine and Clinical Laboratories
Memorial Sloan-Kettering Cancer Center
New York, New York

Mary Jane Massie, MD

Professor of Clinical Psychiatry
Weill Medical College of Cornell University
Attending Psychiatrist
Department of Psychiatry and Behavioral Science
Memorial Sloan-Kettering Cancer Center
New York, New York

Ellen T. Matloff, MS

Associate Research Scientist
Department of Genetics
Yale University School of Medicine
Yale Cancer Center
New Haven, Connecticut

Peter M. Mauch, MD

Professor of Radiation Oncology
Harvard Medical School
Brigham and Women's Hospital
Boston, Massachusetts

Susan Taylor Mayne, PhD, FACS

Associate Professor of Epidemiology and Public Health
Yale University School of Medicine
Yale Cancer Center
New Haven, Connecticut

Minesh P. Mehta, MD

Associate Professor
Interim Chair
Department of Human Oncology
University of Wisconsin Medical School
Madison, Wisconsin

Marvin L. Meistrich, PhD

Professor of Experimental Radiation Oncology
The University of Texas M. D. Anderson
Cancer Center
Houston, Texas

Beth E. Meyerowitz, PhD

Professor of Psychology
University of Southern California School of Medicine
Los Angeles, California

James W. Mier, MD

Associate Professor of Medicine
Harvard Medical School
Beth Israel Deaconess Medical Center
Boston, Massachusetts

Susan D. Miller, PhD

Assistant Professor of Otolaryngology
Georgetown University School of Medicine
Washington, DC

John D. Minna, MD

Professor of Internal Medicine and Pharmacology
Director, Hamon Center for Therapeutic
Oncology Research
University of Texas Southwestern Medical Center
at Dallas
Dallas, Texas

Bruce D. Minsky, MD
Professor and Vice Chair
Department of Radiation Oncology
Memorial Sloan-Kettering Cancer Center
New York, New York

Jeffrey F. Moley, MD
Associate Professor of Surgery
Washington University School of Medicine
St. Louis, Missouri

Monica Morrow, MD
Professor of Surgery
Northwestern University Medical School
Director, Cancer Department of the American College
of Surgeons
Chicago, Illinois

Augusto C. Mota, MD
Senior Fellow in Medicine
Section of Medical Oncology
Yale University School of Medicine
Yale Cancer Center
New Haven, Connecticut

Robert J. Motzer, MD
Associate Attending Physician
Department of Medicine
Memorial Sloan-Kettering Cancer Center
New York, New York

Franco M. Muggia, MD
The Anne Marnick and David H. Cogan Professor
of Oncology
New York University School of Medicine
New York, New York

Nikhil C. Munshi, MD
Professor of Medicine
University of Arkansas for Medical Sciences
Myeloma and Transplantation Research Center
Little Rock, Arkansas

John Murren, MD
Associate Professor of Medicine
Yale University School of Medicine
New Haven, Connecticut

Dao M. Nguyen, MD
Senior Investigator
Department of Surgery
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Jeffrey A. Norton, MD
Professor of Surgery
University of California, San Francisco, School
of Medicine
Chief of Surgery, San Francisco Veterans Affairs
Medical Center
San Francisco, California

Peter J. O'Dwyer, MB, BCH
Professor of Medicine
Division of Hematology-Oncology
University of Pennsylvania School of Medicine
Philadelphia, Pennsylvania

Edward H. Oldfield, MD
Chief, Surgical Neurology Branch
National Institute for Allergies and Infectious Diseases,
National Institutes of Health
Bethesda, Maryland

C. Kent Osborne, MD
Professor of Medicine and Molecular and
Cellular Biology
Director, Breast Center
Baylor College of Medicine
Houston, Texas

Willem W. Overwijk, MS
Surgery Branch
National Cancer Institute
Bethesda, Maryland

Robert F. Ozols, MD, PhD
Senior Vice President
Division of Medical Science
Fox Chase Cancer Center
Philadelphia, Pennsylvania

Harvey I. Pass, MD
Professor of Surgery and Oncology
Department of Cardiothoracic Surgery
Wayne State University School of Medicine
Karmanos Cancer Institute
Detroit, Michigan

Roy Patchell, MD
Associate Professor of Surgery (Neurosurgery)
University of Kentucky College of Medicine
Chief of Neuro-Oncology
Division of Neurosurgery
University of Kentucky Medical Center
Lexington, Kentucky

Peter L. Perrotta, MD
Assistant Professor of Pathology
SUNY at Stony Brook School of Medicine
University Hospital
Stony Brook, New York

Jeanne A. Petrek, MD
Attending Surgeon
Memorial Sloan-Kettering Cancer Center
New York, New York

Eric M. Poeschla, MD
Molecular Medicine Program and Division of
Infectious Diseases
Mayo Medical School
Mayo Clinic
Rochester, Minnesota

Arthur S. Polans, PhD
Jules and Doris Stein RPB Professor
Department of Ophthalmology
and Visual Sciences
University of Wisconsin Medical School
Madison, Wisconsin

Martin G. Pomper, MD, PhD

Assistant Professor of Radiology
Division of Neuroradiology
Johns Hopkins University School of Medicine
Baltimore, Maryland

Amanda Psyrri, MD

Postdoctoral Fellow in Medical Oncology
Yale University School of Medicine
Yale-New Haven Hospital
New Haven, Connecticut

Joe B. Putnam, Jr., MD

Professor of Surgery
Department of Thoracic and Cardiovascular Surgery
The University of Texas M. D. Anderson Cancer Center
Houston, Texas

Mazin B. Qumsiyeh, PhD, FABMG

Associate Professor of Genetics
Yale University School of Medicine
New Haven, Connecticut

Mark J. Ratain, MD

Professor of Medicine
Chair, Committee on Clinical Pharmacology
University of Chicago Pritzker School of Medicine
Chicago, Illinois

Dianne M. Reeves, RN

Senior Research Nurse
Medicine Branch, Division of Clinical Sciences
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Nicholas P. Restifo, MD

Surgery Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Stanley R. Riddell, MD

Professor of Medicine
University of Washington School of Medicine
Member, Fred Hutchinson Cancer Research Center
Program in Immunology
Seattle, Washington

Barbara K. Rimer, DrPH

Director, Division of Cancer Control and Population Sciences
National Cancer Institute, National Institutes
of Health
Rockville, Maryland

Brad Rodu, DDS

Professor of Pathology
University of Alabama School of Medicine
Birmingham, Alabama

Steven A. Rosenberg, MD, PhD

Chief of Surgery
National Cancer Institute, National Institutes of Health
Professor of Surgery
Uniformed Services University of the Health Sciences School
of Medicine
Bethesda, Maryland
Professor of Surgery
George Washington University School of Medicine
Washington, DC

Kenneth Rosenzweig, MD

Assistant Member
Department of Radiation Oncology
Memorial Sloan-Kettering Cancer Center
New York, New York

Andrew J. Roth, MD

Assistant Attending
Department of Psychiatry and Behavioral Sciences
Memorial Sloan-Kettering Cancer Center
New York, New York

Eric K. Rowinsky, MD, FACP

Director, Clinical Research
Institute for Drug Development
Cancer Therapy and Research Center
San Antonio, Texas

José A. Sahel, MD

Professor of Ophthalmology
Clinique Opthalmologique
Hôpitaux Universitaires de Strasbourg
Strausbourg, France

Kapaettu Satyamoorthy, PhD

Senior Scientist
Department of Molecular and Cellular Oncology
The Wistar Institute
Philadelphia, Pennsylvania

Stimson P. Schantz, MD

Professor of Otolaryngology
The New York Eye and Ear Infirmary
New York, New York

David A. Scheinberg, MD, PhD

Professor of Medicine
Member
Department of Medical Oncology
Memorial Sloan-Kettering Cancer Center
New York, New York

Peter B. Schiff, MD, PhD

Professor and Chair of Radiation Oncology
Columbia University College of Physicians
and Surgeons
New York-Presbyterian Hospital
New York, New York

Joellen Schildkraut, PhD

Associate Professor of Community and
Family Medicine
Duke University Medical Center
Durham, North Carolina

John T. Schiller, MD

Division of Basic Sciences
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

David S. Schrump, MD

Head, Thoracic Oncology
Department of Surgery
National Cancer Institute, National Institute
of Health
Bethesda, Maryland

Peter E. Schwartz, MD

Professor of Obstetrics and Gynecology
Yale University School of Medicine
New Haven, Connecticut

Douglas J. Schwartzentruber, MD

Senior Investigator
Surgery Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Brahm H. Segal, MD

Assistant Professor of Medicine
Department of Infectious Diseases and Allergy,
Immunology, and Rheumatology
University at Buffalo School of
Medicine and Biomedical Sciences
Buffalo, New York

Claudia A. Seipp, RN, OCN

Oncology Nurse Clinician
Surgery Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Yoshitaka Sekido, MD, PhD

Assistant Professor of Clinical Preventive Medicine
Nagoya University School of Medicine
Nagoya, Aichi, Japan

Stuart Seropian, MD

Assistant Professor of Medicine
Department of Internal Medicine, Section of
Medical Oncology
Yale University School of Medicine
New Haven, Connecticut

Roy B. Sessions, MD, FACS

Professor of Otolaryngology/Head and
Neck Surgery
Albert Einstein College of Medicine of
Yeshiva University
Beth Israel Medical Center
New York, New York

Jay Shah, MD

Rehabilitation Medicine Department
National Institutes of Health
Bethesda, Maryland

Robert C. Shamberger, MD

Professor of Surgery
Department of Pediatric Surgery
Harvard Medical School
Children's Hospital
Boston, Massachusetts

Joel Sheinfeld, MD

Associate Member
Department of Urology
Memorial Sloan-Kettering Cancer Center
New York, New York

Richard M. Sherry, MD

Senior Investigator
Surgery Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Peter G. Shields, MD

Professor of Medicine and Oncology
Cancer Genetics and Epidemiology Program
Georgetown University School of Medicine
Vincent T. Lombardi Cancer Research Center
Washington, DC

William U. Shipley, MD, FACP

Professor of Radiation Oncology
Harvard Medical School
Chair Genitourinary Oncology
Department of Radiation Oncology
Massachusetts General Hospital
Boston, Massachusetts

David Sidransky, BS, MD

Professor of Otolaryngology, Head and
Neck Surgery
Johns Hopkins University School
of Medicine
Baltimore, Maryland

Mark Siegler, MD

Lindy Bergman Distinguished Service Professor
of Medicine
Director, MacLean Center for Clinical Medical Ethics
University of Chicago Pritzker School of Medicine
Chicago, Illinois

Richard Simon, DSc

Chief, Biometric Research Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Stephen X. Skapek, MD

Assistant Member
Department of Hematology-Oncology (Molecular
Therapeutics)
St. Jude Children's Research Hospital
Memphis, Tennessee

Monica Skarulis, MD

Senior Clinical Investigator
National Institute of Diabetes and Digestive and
Kidney Diseases, National Institutes of Health
Bethesda, Maryland

John M. Skibber, MD

Associate Professor of Surgery
Department of Surgical Oncology
The University of Texas M. D. Anderson
Cancer Center
Houston, Texas

Alfred R. Smith, PhD

Professor of Radiation Oncology
Harvard Medical School
Massachusetts General Hospital
Boston, Massachusetts

XX Contributing Authors

J. Stanley Smith, MD

Professor of Surgery
Pennsylvania State University College of Medicine
Milton S. Hershey Medical Center
Hershey, Pennsylvania

Phyllis M. Smith, MS, LCSW

Senior Clinical Social Worker
Department of Social Work
Yale-New Haven Medical Center
New Haven, Connecticut

Edward L. Snyder, MD

Professor of Laboratory Medicine
Yale University School of Medicine
Director, Blood Bank
Yale-New Haven Hospital
New Haven, Connecticut

Wiley W. Souba, MD, ScD, MBA

Waldhausen Professor and Chair
Department of Surgery
Pennsylvania State University College
of Medicine
Hershey, Pennsylvania

Ira J. Spiro, MD, PhD

Department of Radiation Oncology
Harvard Medical School
Massachusetts General Hospital
Boston, Massachusetts

Laurel J. Steinherz, MD

Associate Professor of Clinical Pediatrics
Weill Medical College of Cornell University
Associate Member
Memorial Sloan-Kettering Cancer Center
Director of Pediatric Cardiology
Memorial Hospital for Cancer and Allied Diseases
New York, New York

William G. Stetler-Stevenson, MD, PhD

Senior Investigator, Chief, Extracellular Matrix
Pathology Section
Laboratory of Pathology
Division of Clinical Sciences
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

James P. Stevenson, MD

Assistant Professor of Medicine
Department of Medical Oncology
University of Pennsylvania School of Medicine
Philadelphia, Pennsylvania

Clinton F. Stewart, PharmD

Associate Member
Department of Pharmaceutical Sciences
St. Jude Children's Research Hospital
Memphis, Tennessee

Diane E. Stover, MD

Professor of Clinical Medicine
Weill Medical College of Cornell University
Chief, Pulmonary Medicine
Head, General Medicine
Memorial Sloan-Kettering Cancer Center
New York, New York

Mario Sznol, MD

Vice President
Department of Clinical Affairs
Vion Pharmaceuticals
New Haven, Connecticut

Moshe Talpaz, MD

Professor of Medicine
Department of Bioimmunotherapy
The University of Texas M. D. Anderson Cancer Center
Houston, Texas

Nancy J. Tarbell, MD

Professor of Radiation Oncology
Harvard Medical School
Massachusetts General Hospital
Boston, Massachusetts

Joel E. Tepper, MD

Professor and Chair
Department of Radiation Oncology
University of North Carolina at Chapel Hill School of Medicine
Chapel Hill, North Carolina

Joseph R. Testa, PhD

Senior Member
Director, Human Genetics Program
Fox Chase Cancer Center
Philadelphia, Pennsylvania

James T. Thigpen, MD

Professor of Medicine
University of Mississippi School of Medicine
University of Mississippi Medical Center
Jackson, Mississippi

Charles R. Thomas, Jr., MD

Associate Professor and Vice Chair
Department of Radiation Oncology
Adjunct Associate Professor of Medical Oncology
University of Texas Health Science Center
at San Antonio
San Antonio Cancer Institute
San Antonio, Texas

Michael J. Thun, MD, MS

Vice President
Department of Epidemiology and
Surveillance Research
American Cancer Society
Atlanta, Georgia

Anthony W. Tolcher, MD, FRCP(C)

Assistant Clinical Professor of Medicine
University of Texas Health Science Center
at San Antonio
San Antonio, Texas

Lols B. Travis, MD, ScD

Senior Investigator
Division of Cancer Epidemiology and Genetics
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Guldo Tricot, MD, PhD

Director, Myeloma Treatment and Transplantation
Research Center
University of Arkansas for Medical Sciences
Little Rock, Arkansas

Margaret A. Tucker, MD

Chief, Genetic Epidemiology Branch
Division of Cancer Epidemiology and Genetics
National Cancer Institute, National Institutes
of Health
Rockville, Maryland

Robert L. Ullrich, PhD

Professor of Radiation Oncology
University of Texas Medical School
at Galveston
Galveston, Texas

Bruce A. Urban, MD

Assistant Professor of Radiology
Johns Hopkins University School
of Medicine
Johns Hopkins Hospital
Baltimore, Maryland

Flora E. van Leeuwen, PhD

Professor of Cancer Epidemiology
Department of Epidemiology
The Netherlands Cancer Institute
Amsterdam, The Netherlands

Rena Vassilopoulou-Sellin, MD, FACP, FACE

Professor of Medicine
Internist
Department of Internal Medicine Specialties/
Endocrinology
The University of Texas M. D. Anderson
Cancer Center
Houston, Texas

Everett E. Vokes, MD

Duchossois Professor of Medicine and
Radiation Oncology
Director, Section of Hematology/Oncology
University of Chicago Pritzker School
of Medicine
Chicago, Illinois

Margaret von Mehren, MD

Associate Member
Department of Medical Oncology
Fox Chase Cancer Center
Philadelphia, Pennsylvania

Thomas J. Walsh, MD

Senior Investigator
Pediatric Oncology Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

McClellan M. Walther, MD

Department of Urology Oncology
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Raymond P. Warrell, Jr., MD

President and Chief Executive Officer
Genta Incorporated
Berkeley Heights, New Jersey

Louis M. Weiner, MD

Chair and Professor of Medicine
Department of Medical Oncology
Fox Chase Cancer Center
Philadelphia, Pennsylvania

Howard J. Weinstein, MD

Professor of Pediatrics
Pediatric Hematology-Oncology
Harvard Medical School
MassGeneral Hospital for Children
Boston, Massachusetts

Mark Weiss, MD

Assistant Professor of Medicine
Memorial Sloan-Kettering Cancer Center
New York, New York

Raymond B. Weiss, MD

Clinical Professor of Medicine
Georgetown University School of Medicine
Washington, D.C.

Samuel A. Wells, Jr., MD

Professor of Surgery
Washington University School of Medicine
St. Louis, Missouri

Patrick Y. Wen, MD

Associate Professor of Neurology
Harvard Medical School
Center for Neuro-Oncology
Dana-Farber Cancer Institute
Division of Neuro-Oncology
Brigham and Women's Hospital
Boston, Massachusetts

Jeffrey D. White, MD

Director
Office of Cancer Complementary and
Alternative Medicine
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Christopher G. Willett, MD

Professor of Radiation Oncology
Harvard Medical School
Massachusetts General Hospital
Boston, Massachusetts

Walter C. Willett, MD

Department of Nutrition
Harvard School of Public Health
Boston, Massachusetts

Lynn D. Wilson, MD, MPH

Associate Professor of Therapeutic Radiology
Yale University School of Medicine
New Haven, Connecticut

Eric P. Winer, MD

Associate Professor of Medicine
Department of Adult Oncology
Dana-Farber Cancer Institute
Boston, Massachusetts

Flossie Wong-Staal, PhD

Florence Seeley Riford Professor in AIDS Research
Department of Medicine and Biology
University of California, San Diego, School
of Medicine
La Jolla, California

John R. Wunderlich, MD

Surgery Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Joachim Yahalom, MD

Professor of Radiation Oncology in Medicine
Weill Medical College of Cornell University
Attending and Member
Memorial Sloan-Kettering Cancer Center
New York, New York

James C. Yang, MD

Senior Investigator
Surgery Branch
National Cancer Institute, National Institutes
of Health
Bethesda, Maryland

Robert Yarchoan, MD

Chief, HIV and AIDS Malignancy Branch
Division of Clinical Sciences
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Stuart H. Yuspa, MD

Laboratory Chief
Laboratory of Cellular Carcinogenesis and Tumor Promotion
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Berton Zbar

Chief, Laboratory of Immunology
Division of Basic Sciences
National Cancer Institute, National Institutes of Health
Bethesda, Maryland

Michael J. Zelefsky, MD

Associate Professor of Radiation Oncology
Director of Brachytherapy
Memorial Sloan-Kettering Cancer Center
New York, New York

Sandra S. Zinkel, MD, PhD

Research Fellow in Cancer Immunology and AIDS
Dana-Farber Cancer Institute
Boston, Massachusetts

EXHIBIT G

Your continued donations keep Wikipedia running!

RET proto-oncogene

From Wikipedia, the free encyclopedia

The **RET proto-oncogene** encodes a receptor tyrosine kinase for members of the glial cell line-derived neurotrophic factor family of extracellular signalling molecules.^[1] *RET* loss of function mutations are associated with the development of Hirschsprung's disease, while gain of function mutations are associated with the development of various types of human cancer, including medullar thyroid carcinoma and multiple endocrine neoplasias type 2A and type 2B.

Contents
<div><ul style="list-style-type: none">1 Structure2 Kinase activation3 Role of RET signalling during development4 References5 External links</div>

Structure

RET is an abbreviation for "rearranged during transfection", as the DNA sequence of this gene was originally found to be rearranged within a 3T3 fibroblast cell line following its transfection with DNA taken from human lymphoma cells.^[2] The human gene *RET* is localized to chromosome 10 (10q11.2) and contains 21 exons.^[3]

The natural alternate splicing of the *RET* gene results in the production of 3 different isoforms of the protein RET. RET51, RET43 and RET9 contain 51, 43 and 9 amino acids in their C-terminal tail respectively.^[4] The biological roles of isoforms RET51 and RET9 are the most well studied *in-vivo* as these are the most common isoforms in which RET occurs.

Common to each isoform is a domain structure. Each protein is divided into three domains: an N-terminal extracellular domain with four

	ret proto-oncogene
Identifiers	
Symbol	RET
Alt. Symbols	HSCR1, MEN2A, MTC1, MEN2B
Entrez	5979 (http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene&cmd=retrieve&dopt=default&list_uids=5979&rn=1)
HUGO	9967 (http://www.genenames.org/data/hgnc_data.php?hgnc_id=9967)
OMIM	164761 (http://www.ncbi.nlm.nih.gov/entrez/dispmim.cgi?id=164761&rn=1)
RefSeq	NM_020975 (http://genome.cse.ucsc.edu/cgi-bin/hgGene?org=Human&hgg_gene=NM_020975&rn=1)
UniProt	P07949 (http://www.expasy.org/cgi-bin/niceprot.pl?P07949)
Other data	
Locus	Chr. 10 <i>q11.2</i> (http://www.ncbi.nlm.nih.gov/Omim/getmap.cgi?chromosome=10q11.2)

cadherin-like repeats and a cysteine-rich region, a hydrophobic transmembrane domain and a cytoplasmic tyrosine kinase domain, which is split by an insertion of 27 amino acids. Within the cytoplasmic tyrosine kinase domain, there are 16 tyrosines (Tyr) in RET9 and 18 in RET51. Tyr1090 and Tyr1096 are present only in the RET51 isoform.^[5]


The extracellular domain of RET contains nine N-glycosylation sites. The fully glycosylated RET protein is reported to have a molecular weight of 170 kDa although it is not clear to which isoform this molecular weight relates.^[6]

Kinase activation

RET is the receptor for members of the glial cell line-derived neurotrophic factor (GDNF) family of extracellular signalling molecules or ligands (GFLs).^[7]

In order to activate RET GFLs first need to form a complex with a glycosylphosphatidylinositol (GPI)-anchored co-receptor. The co-receptors themselves are classified as members of the GDNF receptor- α (GFR α) protein family. Different members of the GFR α family (GFR α 1-GFR α 4) exhibit a specific binding activity for a specific GFLs.^[8] Upon GFL-GFR α complex formation, the complex then brings together two molecules of RET, triggering trans-autophosphorylation of specific tyrosine residues within the tyrosine kinase domain of each RET molecule. Tyr900 and Tyr905 within the activation loop (A-loop) of the kinase domain have been shown to be autophosphorylation sites by mass spectrometry.^[9] Phosphorylation of Tyr905 stabilizes the active conformation of the kinase which in turn results in the autophosphorylation of other tyrosine residues mainly located in the C-terminal tail region of the molecule.^[5]

The structure shown below was taken from the protein data bank code 2IVT (<http://www.pdb.org/pdb/explore.do?structureId=2IVT>).^[1] The structure is that of a dimer formed between two protein molecules each spanning from amino acids 703-1012 of the RET molecule, covering RETs intracellular tyrosine kinase domain. One protein molecule, molecule A is shown in yellow and the other, molecule B in grey. The activation loop is coloured purple and selected tyrosine residues in green. Part of the activation loop from molecule B is absent.



Phosphorylation of Tyr981 and the additional tyrosines Tyr1015, Tyr1062 and Tyr1096 not covered by the above structure, have been shown to be important to the initiation of intracellular signal transduction processes.

Role of RET signalling during development

Mice deficient in GDNF, GFR α 1 or the RET protein itself exhibit severe defects in kidney and enteric nervous system development. This implicates RET signal transduction as key to the development of normal kidneys and the enteric nervous system.^[5]

References

- ^{a b} Knowles PP, Murray-Rust J. *et al* (2006). "Structure and chemical inhibition of the RET tyrosine kinase domain". *J. Biol. Chem.* **281** (44): 33577-33587. PMID 16928683 (http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&dopt=Abstract&list_uids=16928683).
- ^a Takahashi M, Ritz J, Cooper GM. (1985). "Activation of a novel human transforming gene, ret, by DNA rearrangement." *Cell* **42** (2): 581-588. PMID 2992805 (http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&dopt=Abstract&list_uids=2992805).
- ^a Ceccherini I, Bocciardi R. *et al* (1993). "Exon structure and flanking intronic sequences of the human RET proto-oncogene". *Biochem. Biophys. Res. Commun.* **196** (3): 1288-1295. PMID 7902707 (http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&dopt=Abstract&list_uids=7902707).
- ^a Myers SM, Eng C. *et al* (1995). "Characterization of RET proto-oncogene 3' splicing variants and polyadenylation sites: a novel C-

terminus for RET". *Oncogene* **11** (10): 2039-2045. PMID 7478523 (http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&dopt=Abstract&list_uids=7478523).

5. ^{a b c} Arighi E, Borrello MG, Sariola H. (2005). "RET tyrosine kinase signaling in development and cancer". *Cytokine Growth Factor Rev.* **16** (4-5): 441-467. PMID 15982921 (http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&dopt=Abstract&list_uids=15982921).
6. ^a Takahashi M, Asai N. *et al* (1993). "Characterization of the ret proto-oncogene products expressed in mouse L cells." *Oncogene* **8** (11): 2925-2929. PMID 8414495 (http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&dopt=Abstract&list_uids=8414495).
7. ^a Baloh RH, Enomoto H. *et al* (2000). "The GDNF family ligands and receptors - implications for neural development". *Curr. Opin. Neurobiol.* **10** (1): 103-110. PMID 10679429 (http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&dopt=Abstract&list_uids=10679429).
8. ^a Airaksinen MS, Titevsky A, Saarna M. (1999). "GDNF family neurotrophic factor signaling: four masters, one servant?". *Mol. Cell Neurosci.* **13** (5): 313-325. PMID 10356294 (http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&dopt=Abstract&list_uids=10356294).
9. ^a Kawamoto Y, Takeda K. *et al* (2004). "Identification of RET autophosphorylation sites by mass spectrometry". *J. Biol. Chem.* **279** (14): 14213-14224. PMID 14711813 (http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&dopt=Abstract&list_uids=14711813).

External links

- MeSH *ret+Proto-Oncogene+Proteins* (http://www.nlm.nih.gov/cgi/mesh/2007/MB_cgi?mode=&term=ret+Proto-Oncogene+Proteins)

Retrieved from "http://en.wikipedia.org/wiki/RET_proto-oncogene"

Categories: Protein pages needing a picture | Genes on chromosome 10 | Tyrosine kinase receptors

- ▣ This page was last modified 22:21, 11 June 2007.
 - ▣ All text is available under the terms of the GNU Free Documentation License. (See **Copyrights** for details.)
- Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a U.S. registered 501(c)(3) tax-deductible nonprofit charity.